ARTHAM
RESOURCE MATERIAL

SCHOOL OF EDUCATORS
SOE
Learn / Unlearn / Relearn

COMPETENCY-BASED QUESTION BANK

WITH ANSWER KEY
& STRUCTURED EXPLANATION

CLASS 11
MATHEMATICS

FEATURES

1 Comprehensive Coverage
Includes detailed question banks for Class 11 & 12 subjects PCMB

2 Skill-based Learning
Develops critical thinking and problem-solving skills essential for cracking tough exams.

3 Score Boosting
Helps to score maximum marks in CBSE exams and increases competitive exam success potential..

4 Answer Key & Structured Explanations
Clear, well-structured explanations and step-by-step solutions to enhance understanding.

5 Exam Pattern Aligned
Questions modeled after the latest exam patterns to ensure students are well-prepared.

6 Competitive Exam Focus
Specially designed to help students excel in IIT, NEET, JEE, CUET, and other competitive exams.

7 NCERT Integration
Questions and answers are based on the NCERT syllabus, ensuring relevance for both CBSE board exams and entrance tests.

ARTHAM RESOURCES

Class: 11,12 Mathematics
Competency-based Question Bank with Answer Key
& Structured Explanation

LINEAR INEQUALITIES

1. If $3^x + 2^{2x} \geq 5^x$, then the solution set for x is
   a) $(-\infty, 2]$ b) $[2, \infty)$ c) $[0, 2]$ d) $\{2\}$
2. $x^2 - 3|x| + 2 < 0$, then x belongs to
   a) $(1, 2)$ b) $(-2, -1)$ c) $(-2, -1) \cup (1,2)$ d) $(-3, 5)$
3. Solution of $2^x + 2^{|x|} \geq 2\sqrt{2}$ is
   a) $(-\infty, \log_2(\sqrt{2} + 1))$ b) $(0,8)$
   c) $\left(\frac{1}{2}, \log_2(\sqrt{2} - 1)\right)$ d) $(-\infty, \log_2(\sqrt{2} - 1)) \cup \left[\frac{1}{2}, \infty\right)$
4. If $x_1, x_2, \ldots, x_n$ are real numbers, then the largest value of the expression $\sin x_1 \cos x_2 + \sin x_2 \cos x_3 + \cdots + \sin x_n \cos x_1$ is
   a) n b) $\frac{n}{2}$ c) $\frac{n}{4}$ d) $\sqrt{n^2 - 1}$
5. If $a < b$, then the solution $x^2 + (a + b)x + ab < 0$ is given by
   a) (a, b) b) $(-\infty, a) \cup (b, \infty)$ c) $(-b, -a)$ d) $(-\infty, -b) \cup (-a, \infty)$
6. If $\log(x^3 + y^3) - \log_{10}(x^2 + y^2 - xy) \leq 2$, then the maximum value of xy, for all $x \geq 0, y \geq 0$ is
   a) 2500 b) 3000 c) 1200 d) 3500
7. If $3^{x/2} + 2^x > 25$, then the solution set is
   a) R b) $(2, \infty)$ c) $(4, \infty)$ d) None of these
8. If $ab = 4 (a, b \in R^+)$, then
   a) $a + b \leq 4$ b) $a + b = 4$ c) $a + b \geq 4$ d) None of these
9. Let $P_n(x) = 1 + 2x + 3x^2 + \cdots + (n + 1)x^n$ be a polynomial such that n is even. Then, the number of real roots of $P_n(x)$, is
   a) 0 b) n c) 1 d) None of these
10. $(x - 1)(x^2 - 5x + 7) < (x - 1)$, then x belongs to
    a) $(1,2) \cup (3, \infty)$ b) $(2, 3)$ c) $(-\infty, 1) \cup (2,3)$ d) None of these
11. If $x = \log_{2^2} 2 + \log_{2^3} 2^2 + \log_{2^4} 2^3 + \ldots + \log_{2^{n+1}} 2^n$, then
    a) $x \geq \left(\frac{1}{n+1}\right)^{1/n}$ b) $x \geq n\left(\frac{1}{n+1}\right)^{1/n}$ c) $x \geq \left(\frac{n}{n+1}\right)^{1/n}$ d) None of these
12. The number of real solutions (x, y, z, t) of simultaneous equations
    $2y = \frac{11}{x} + x, 2z = \frac{11}{y} + y, 2t = \frac{11}{z} + z, 2x = \frac{11}{t} + t$, is
    a) 0 b) 1 c) 2 d) 4
13. The solution set contained in R of the inequation $3^x + 3^{1-x} - 4 < 0$, is
    a) $(1, 3)$ b) $(0, 1)$ c) $(1, 2)$ d) $(0, 2)$
14. The range of ab if $|a| \leq 1$ and $a + b = 1, (a, b \in R)$, is
    a) $[0, 1/4]$ b) $[-2, 1/4]$ c) $[1/4, 2]$ d) $[0, 2]$
15. If $\sqrt{9x^2 + 6x + 1} < (2 - x)$, then
    a) $x \in \left(-\frac{3}{2}, \frac{1}{4}\right)$ b) $x \in \left(-\frac{3}{2}, \frac{1}{4}\right)$ c) $x \in \left[-\frac{3}{2}, \frac{1}{4}\right]$ d) $x < \frac{1}{4}$
16. If $5^x + \left(2\sqrt{3}\right)^{2x} \geq 13^x$, then the solution set for x is
    a) $[2, \infty)$ b) $\{2\}$ c) $(-\infty, 2]$ d) $[0, 2]$
17. Solution set of inequality $\log_e \frac{x-2}{x-3}$ is
    a) $(2, \infty)$ b) $(-\infty, 2)$ c) $(-\infty, \infty)$ d) $(3, \infty)$
18. If $3 < 3t - 18 \leq 18$, then which one of the following is true?
    a) $15 \leq 2t + 1 \leq 20$ b) $8 \leq t < 12$ c) $8 \leq t + 1 \leq 13$ d) $21 \leq 3t \leq 24$
19. Let $f(x) = ax^2 + bx + c$ and $f(-1) < 1, f(1) > -1, f(3) < -4$ and $a \neq 0$, then
    a) $a > 0$ b) $a < 0$
    c) Sign of a cannot be determined d) None of the above

20. The set of admissible values of x such that $\frac{2x+3}{2x-9} < 0$ is

a) $\left(-\infty, -\frac{3}{2}\right) \cup \left(\frac{9}{2}, \infty\right)$ b) $(-\infty, 0) \cup \left(\frac{9}{2}, \infty\right)$ c) $\left(-\frac{3}{2}, 0\right)$ d) $\left(-\frac{3}{2}, \frac{9}{2}\right)$

21. The number of irrational solutions of the equation $\sqrt{x^2 + \sqrt{x^2 + 11}} + \sqrt{x^2 - \sqrt{x^2 + 11}} = 4$, is

a) 0 b) 2 c) 4 d) 11

22. The number of solutions of the equation $\log_{x-3}(x^3 - 3x^2 - 4x + 8) = 3$, is

a) 1 b) 2 c) 3 d) 4

23. The number of real solutions of the equation $\log_{0.5} x = |x|$, is

a) 1 b) 0 c) 2 d) None of these

24. The number of complex roots of the equation $x^4 - 4x - 1 = 0$, is

a) 3 b) 2 c) 1 d) 0

25. If $\sin^x \alpha + \cos^x \alpha \geq 1, 0 < \alpha < \frac{\pi}{2}$,then

a) $x \in (2, \infty)$ b) $x \in (-\infty, 2]$ c) $x \in [-1, 1]$ d) None of these

26. Consider the following statements:

1. $\frac{x}{1+x^2} < \tan^{-1} x < x; x > 0$

2. If $0 \leq x < \frac{\pi}{2}, \sin x + \tan x - 3x \geq 0$

Which of these is/are correct?

a) Only (1) b) Only (2) c) (1) and (2) d) None of these

27. The number of solutions of the equation $2\cos(e^x) = 3^x + 3^{-x}$, is

a) 0 b) 1 c) 2 d) None of these

28. The number of real solutions of the equation $1 - x = [\cos x]$, is

a) 1 b) 2 c) 3 d) None of these

29. Non- negative real numbers such that $a_1 + a_2 + \ldots + a_n = p$ and $q = \sum_{i<j} a_i a_j$,then

a) $q \leq \frac{1}{2}p^2$ b) $q > \frac{1}{4}p^2$ c) $q < \frac{p}{2}$ d) $q > \frac{p^2}{2}$

30. If $(\sin a)^x + (\cos a)^x \geq 1, 0 < a < \frac{\pi}{2}$, then

a) $x \in [2, \infty)$ b) $x \in (-\infty, 2]$ c) $x \in [-1, 1]$ d) None of these

31. If $x^2 + 2ax + 10 - 3a > 0$ for all $x \in R$, then

a) $-5 < a < 2$ b) $a < -5$ c) $a > 5$ d) $2 < a < 5$

32. The least integer satisfying$49.4 - \left(\frac{27-x}{10}\right) < 47.4 - \left(\frac{27-9x}{10}\right)$, is

a) 2 b) 3 c) 4 d) None of these

33. For positive real number a, b, c which one of the following holds?

a) $a^2 + b^2 + c^2 \geq bc + ca + ab$ b) $(b + c)(c + a)(a + b) \leq 8abc$

c) $\frac{a}{b} + \frac{b}{c} + \frac{c}{a} \leq 3$ d) $a^3 + b^3 + c^3 \leq 3\,abc$

34. The least perimeter of a cyclic quadrilateral of given area A square units is

a) $\sqrt{A}$ b) $2\sqrt{A}$ c) $3\sqrt{A}$ d) $4\sqrt{A}$

35. The number of solutions of $[\sin x + \cos x] = 3 + [-\sin x] + [-\cos x]$ in the internal $[0, 2\pi]$ is (where [.] denotes the greatest integer function)

a) 0 b) 4 c) Infinite d) 1

36. The number of solutions of $3^{|x|} = \left|2 - |x|\right|$ is

a) 0 b) 2 c) 4 d) Infinite

37. If C is an obtuse angle in tringle, then

a) $\tan A \tan B < 1$ b) $\tan A \tan B > 1$ c) $\tan A \tan B = 1$ d) None of these

38. If x, y, z are three real numbers such that $x + y + z = 4$ and $x^2 + y^2 + z^2 = 6$, then the exhaustive set of values of x, is

a) $[2/3, 2]$ b) $[0, 2/3]$ c) $[0, 2]$ d) $[-1/3, 2/3]$

39. The number of roots of the equation $[\sin^{-1} x] = x - [x]$, is

a) 0 b) 1 c) 2 d) None of these

40. If $3^{x/2} + 2^x > 25$, then

a) $x \in [4, \infty)$ b) $(4, \infty)$ c) $x \in (-\infty, 4]$ d) $x \in [0, 4]$

41. The number of real solutions of the equation $27^{1/x} + 12^{1/x} = 2.8^{1/x}$, is

a) 1 b) 2 c) 0 d) Infinite

42. If roots of the equation $x^4 - 8x^3 + bx^2 + cx + 16 = 0$ are positive then

a) $b = 8 = c$ b) $b = -24, c = -32$ c) $b = 24, c = -32$ d) $b = 24, c = 32$

43. If $3<|x|<6$, then x belongs to

a) $(-6, -3) \cup (3,6)$ b) $(-6, 6)$ c) $(-3, -3) \cup (3,6)$ d) None of these

44. If a, b are distinct positive real numbers, then which one of the following is true?

a) $a^4 + b^4 > a^3b + ab^3$ b) $a^4 + b^4 < a^3b + ab^3$ c) $a^3 + b^3 < a^2b + ab^2$ d) None of these

45. The solution of the inequation $4^{-x+0.5} - 7 \cdot 2^{-x} < 4, x \in R$ is

a) $(-2, \infty)$ b) $(2, \infty)$ c) $\left(2, \frac{7}{2}\right)$ d) None of tnese

46. Suppose a, b and c are real numbers such that $\frac{a}{b} > 1$ and $\frac{a}{c} < 0$. Which one of the following is true?

a) $a + b - c > 0$ b) $a > b$ c) $(a - c)(b - c) > 0$ d) $a + b + c > 0$

47. If a, b, c are positive real numbers such that $a + b + c = p$ then, which of the following is true?

a) $(p - a)(p - b)(p - c) \geq \frac{1}{27}p^3$

b) $(p - a)(p - b)(p - c) \geq 8abc$

c) $\frac{bc}{a} + \frac{ca}{b} + \frac{ab}{c} \geq p$

d) None of these

48. The number of solutions of the equation $\frac{(1+e^{x^2})\sqrt{1+x^2}}{\sqrt{1+x^4}-x^2} = 1 + \cos x$, is

a) 1 b) 2 c) 3 d) 4

49. Let n be an odd integer such that the polynomial $P_n(x) = 1 + 2x + 3x^2 + \ldots\ldots + (n + 1)x^n$ has exactly one real root. This real root α satisfies

a) $-1 < \alpha < 0$ b) $0 < \alpha < 1$ c) $0 \leq \alpha \leq 1$ d) $-1 \leq \alpha \leq 0$

50. Let a, b be integers and $f(x)$ be a polynomial with integer coefficients such that $f(b) - f(a) = 1$. Then, the value of $b - a$, is

a) 1 b) -1 c) $1, -1$ d) None of these

51. Let $y = \sqrt{\frac{(x+1)(x-3)}{(x-2)}}$, then all real values of x for which y takes real values, are

a) $-1 \leq x < 2$ or $x \geq 3$ b) $-1 \leq x < 3$ or $x > 2$ c) $1 \leq x < 2$ *or* $x \geq 3$ d) None of these

52. If $a, b, c > 0$ and if $abc = 1$, then the value of $a + b + c + ab + bc + ca$ lies in the interval

a) $(\infty, -6)$ b) $(-6, 0)$ c) $(0, 6)$ d) $(6, \infty)$

53. The number of real roots of the equation $(\sin 2^x)(\cos 2^x) = \frac{2^x + 2^{-x}}{2}$, is

a) 1 b) 2 c) 3 d) None of these

54. The largest interval for which $x^{12} - x^9 + x^4 - x + 1 > 0$ is

a) $-4 < x \leq 0$ b) $0 < x < 1$ c) $-100 < x < 100$ d) $0 < x < \infty$

55. The number of negative real roots of $x^4 - 4x - 1 = 0$, is

a) 3 b) 2 c) 1 d) 0

56. If $0 < x < \frac{\pi}{2}$, then minimum value of $\frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x}$ is

a) $\sqrt{3}$ b) $\frac{1}{2}$ c) $\frac{1}{3}$ d) 1

57. The number of solutions of $\sqrt{3x^2 + 6x + 7} + \sqrt{5x^2 + 10x + 14} = 4 - 2x - x^2$, is

a) 1 b) 2 c) 3 d) 4

58. The solution set of $||x|-1| < |1 - x|, x \in R$ is

a) $(-1,1)$ b) $(0, \infty)$ c) $(-1, \infty)$ d) None of these

59. The minimum value of $f(x) = |3 - x| + 7$ is

a) 0 b) 6 c) 7 d) 8

60. The solution set of the inequation $\frac{x+11}{x-3} > 0$ is

a) $(-\infty, 11) \cup (3, \infty)$ b) $(-\infty, -10) \cup (2, \infty)$ c) $(-100, -11) \cup (1, \infty)$ d) $(-5, 0) \cup (3, 7)$

61. Solution of $2x - 1 = |x + 7|$ is

a) -2 b) 8 c) $-2, 8$ d) None of these

62. The number of positive real roots of $x^4 - 4x - 1 = 0$, is

a) 3 b) 2 c) 1 d) 0

63. The solution set of the inequality $\log_{\sin\left(\frac{\pi}{3}\right)}(x^2 - 3x + 2) \geq 2$ is

a) $\left(\frac{1}{2}, 2\right)$ b) $\left(1, \frac{5}{2}\right)$ c) $\left[\frac{1}{2}, 1\right) \cup \left(2, \frac{5}{2}\right]$ d) None of these

64. If a, b, c are sides of triangle, then $\frac{(a+b+c)^2}{(ab+bc+ca)}$ always belongs to

a) [1, 2] b) [2, 3] c) [3, 4] d) [4, 5]

65. $(x - 1)(x^2 - 5x + 7) < (x - 1)$, then x belongs to

a) $(1, 2) \cup (3, \infty)$ b) $(-\infty, 1) \cup (2, 3)$ c) 2, 3 d) None of these

66. The set of values of x for which the inequalities $x^2 - 3x - 10 < 0$, $10x - x^2 - 16 > 0$ hold simultaneously, is

a) (-2, 5) b) (2, 8) c) (-2, 8) d) (2, 5)

67. The solution of the inequation $\log_{1/3}(x^2 + x + 1) + 1 < 0$ is

a) $(-\infty, -2) \cup (1, \infty)$ b) $[-1, 2]$ c) $(-2, 1)$ d) $(-\infty, \infty)$

68. The number of values of a for which the system of equation $2^{|x|} + |x| = y + x^2 + a$ and $x^2 + y^2 = 1$ has only one solution where a, x, y are real, is

a) 1

b) 2

c) Finitely many but more than 2

d) Infinitely many

69. The solution set of the inequation $\log_{1/3}(x^2 + x + 1) + 1 > 0$, is

a) $(-\infty, -2) \cup (1, \infty)$ b) $[-1, 2]$ c) $(-2, 1)$ d) R

70. The set of all real numbers satisfying the inequation $x^{(\log_{10} x)^2 - 3(\log_{10} x) + 1} > 1000$, is

a) (0, 1000) b) (1000, ∞) c) (0, 100) d) None of these

71. The set of all real x satisfying the inequality $\frac{3-|x|}{4-|x|} > 0$

a) $[-3, 3] \cup (-\infty, -4) \cup (4, \infty)$ b) $(-\infty, -4) \cup (4, \infty)$

c) $(-\infty, -3) \cup (4, \infty)$ d) $(-\infty, -3) \cup (3, \infty)$

72. For positive real number a, b, c which of the following holds?

a) $a + b + c > 3 \Rightarrow a^2 + b^2 + c^2 > 3$ b) $a^6 + b^6 \leq 12a^2b^2 - 64$

c) $a + b + c = \alpha \Rightarrow \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \leq \frac{9}{\alpha}$ d) None of the above

73. If a_1, a_2, a_3 be any positive real numbers, then which of the following statements is not true

a) $3a_1, a_2, a_3 \leq a_1^3 + a_2^3 + a_3^3$

b) $\frac{a_1}{a_2} + \frac{a_2}{a_3} + \frac{a_3}{a_1} \geq 3$

c) $(a_1 + a_2 + a_3)\left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3}\right) \geq 9$

d) $(a_1 + a_2 + a_3)\left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3}\right)^3 \leq 27$

74. If $x^2 + \frac{1}{x^2} = A$ and $x - \frac{1}{x} = B$, where $x \in R$ and $B > 0$, then the minimum value of $\frac{A}{B}$ is

a) $\sqrt{2}$ b) $2\sqrt{2}$ c) $\sqrt{2} + 2$ d) None of these

75. Let n be an odd positive integer. Then, the number of real roots of the polynomial $P_n(x) = 1 + 2x + 3x^2 + \ldots\ldots + (n+1)x^n$, is

a) 0 b) n c) 1 d) None of these

76. The number of positive integers satisfying the inequality $n + 1_{C_{n-2}} - n + 1_{C_{n-1}} \leq 50$ is

a) 9 b) 8 c) 7 d) 6

77. For $\theta > \pi/3$, the value of $f(\theta) = \sec^2\theta + \cos^2\theta$ always lies in the interval

a) $(0, 2)$ b) $[0, 1]$ c) $(1, 2)$ d) $[2, \infty)$

78. If the product of n positive numbers is n^n, then their sum is

a) A positive integer b) Divisible by n c) Equal to $n + \frac{1}{n}$ d) Never less than n^2

79. $\log_2(x^2 - 3x + 18) < 4$,then x belongs to

a) $(1, 2)$ b) $(2, 16)$ c) $(1, 16)$ d) None of these

80. If $[x]^2 = [x + 6]$, where $[x]$ = the greatest integer less than or equal to x, then x must be such that

a) $x = 3, -2$ b) $x \in [-2, -1)$ c) $x \in [3, 4)$ d) $x \in [-2, -1) \cup [3, 4)$

81. If $a, b, c > 0$, the minimum value of $\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b}$ is

a) 1 b) $\frac{3}{2}$ c) 2 d) $\frac{5}{2}$

82. The number of real solutions of the equation $e^{|x|} - |x| = 0$, is

a) 0 b) 1 c) 2 d) None of these

83. If p, q, r are any real numbers, then

a) $\max(p, q) = \max(p, q, r)$ b) $\min(p, q) = \frac{1}{2}(P + q - |p - q|)$

c) $\max(p, q) < \min(p, q, r)$ d) $\max(p, q) = \frac{1}{2}(p + q - |p - q|)$

84. The number of real solutions of $1 + |e^x - 1| = e^x(e^x - 2)$, is

a) 1 b) 2 c) 3 d) 4

85. The solution set of the inequation$\log_{\sin 2\pi/3}(x^2 - 3x + 2) \geq 2$, is

a) $[1/2, 1)$ b) $(2, 5/2]$ c) $[1/2, 1) \cup (2, 5/2]$ d) $[1/2, 5/2]$

86. If $x^2 + 6x - 27 < 0$ and$x^2 - 3x - 4 < 0$, then

a) $x > 3$ b) $x < 4$ c) $3 < x < 4$ d) $x = \frac{7}{2}$

87. If $x, y \in R$, then $\frac{1}{2}(x + y + |x - y|) = x$ holds iff

a) $x > y$ b) $x < y$ c) $x = y$ d) $x \geq y$

88. The set of all x satisfying the inequality $\frac{4x-1}{3x+1} \geq 1$ is

a) $\left(-\infty, -\frac{1}{3}\right) \cup \left[\frac{1}{4}, \infty\right]$ b) $\left(-\infty, -\frac{2}{3}\right) \cup \left[\frac{5}{4}, \infty\right]$ c) $\left(-\infty, -\frac{1}{3}\right) \cup [2, \infty)$ d) $\left(-\infty, -\frac{2}{3}\right)[4, \infty)$

89. The number of solutions of the inequality

$E = 2^{1/\sin^2\alpha_2} \cdot 3^{1/\sin^2\alpha_3} \ldots n^{1/\sin^2\alpha_n} < n!$

Where $\alpha_i \in (-\pi, 2\pi)$ for $i = 2, 3, \ldots, n$ is

a) 0 b) 2^{n-1} c) 3^{n-1} d) None of these

90. The equation $e^x = x(x + 1), x < 0$ has

a) No real roots set

b) Exactly one real root

c) Two real roots

d) Infinitely many real roots

91. Let $F(x)$ be a function defined by $F(x) = x - [x], 0 \neq x \in R$, where $[x]$ is the greatest integer less than or equal to x. Then, the number of solutions of $F(x) + F\left(\frac{1}{x}\right) = 1$

a) 0 b) Infinite c) 1 d) 2

92. The set of all real numbers satisfying the inequation$2^x + 2^{|x|} \geq 2\sqrt{2}$, is

a) $(1/2, \infty)$

b) $(-\infty, \log_2(\sqrt{2}-1))$

c) $(-\infty, 1/2)$

d) $[1/2, \infty) \cup (-\infty, \log_2(\sqrt{2}-1))$

93. Solution of the inequality $\sin^4\left(\frac{x}{3}\right) + \cos^4\left(\frac{x}{3}\right) > \frac{1}{2}$, is given by

a) R b) $\frac{3n\pi}{2} + \frac{3\pi}{4}$

c) $R - \left\{\left(\frac{3n\pi}{2} + \frac{3\pi}{4}\right), n \in I\right\}$ d) None of these

94. If $(\log_5 x)^2 + (\log_5 x) < 2$, then x belongs to the interval

a) $(1/25, 5)$ b) $(1/5, 1/\sqrt{5})$ c) $(1, \infty)$ d) None of these

95. The number of real roots of the equation $x^2 + x + 3 + 2\sin x = 0$ in the interval $[-\pi, \pi]$, is

a) 2 b) 4 c) 6 d) None of these

96. If $\log_{\sqrt{3}}(\sin x + 2\sqrt{2}\cos x) \geq 2, -2\pi \leq x \leq 2\pi$, then the number of solutions of x is

a) 0 b) ∞ c) 3 d) 4

97. Solution of $x^{(\log_{10} x)^2 - 3\log_{10} x + 1} > 1000$ for $x \in R$, is

a) $(10, \infty)$ b) $(100, \infty)$ c) $(1000, \infty)$ d) $(1, \infty)$

98. The largest interval for which $x^{12} - x^9 + x^4 - x + 1 > 0$ is

a) $-4 < x < 0$ b) $0 < x < 1$ c) $-100 < x < 100$ d) $-\infty < x < \infty$

99. The number of roots of the equation $\sin \pi x = |\log|x||$, is

a) 2 b) 4 c) 5 d) 6

100. The equation $\sqrt{x+1} - \sqrt{x-1} = \sqrt{4x-1}$ has

a) No solution

b) One solution

c) Two solutions

d) More than two solutions

101. If $\log_3 x - \log_x 27 < 2$, then x belongs to the interval

a) $(1/3, 27)$ b) $(1/27, 3)$ c) $(1/9, 9)$ d) None of these

102. The set of all solutions of the inequation $x^2 - 2x + 5 \leq 0$ in R is

a) $R - (-\infty, -5)$ b) $R - (5, \infty)$ c) $\emptyset$ d) $R - (-\infty, -4)$

103. $2^{\sin^2 x} + 2^{\cos^2 x}$ is

a) ≤ 2 b) ≥ 2 c) ≤ 1 d) ≥ 1

104. If $a^2 + b^2 + c^2 = 1$, then $ab + bc + ca$ lies in the interval

a) $\left[-\frac{1}{2}, 1\right]$ b) $\left[0, \frac{1}{2}\right]$ c) $[0, 1]$ d) $[1, 2]$

105. The equation $\sqrt{4x+9} - \sqrt{11x+1} = \sqrt{7x+4}$ has

a) No solution

b) One solution

c) Two solutions

d) More than two solutions

106. $\left|x + \frac{2}{x}\right| < 3$, then x belongs to

a) $(-2, -1) \cup (1, 2)$ b) $(-\infty, -2) \cup (-1, 1) \cup (2, \infty)$

c) $(-2, 2)$ d) $(-3, 3)$

107. If a, b, c are the sides of a tringle, then $\frac{a}{b+c-a} + \frac{b}{c+a-b} + \frac{c}{a+b-c}$ is

a) ≤ 3 b) ≥ 3 c) ≥ 2 d) ≤ 2

108. The minimum value of the sum of the lengths of diagonals of a cyclic quadrilateral of area a^2 square units is

a) $\sqrt{2}a$ b) $2\sqrt{2}a$ c) $2a$ d) None of these

109. $|2x - 3| < |x + 5|$, then x belongs to

a) $(-3,5)$ b) $(5,9)$ c) $\left(-\frac{2}{3},8\right)$ d) $\left(-8,\frac{2}{3}\right)$

110. The number of real roots of the equation $1+a_1x+a_2x^2+\cdots+a_nx^n=0$, where $|x|<\frac{1}{3}$ and $|a_n|<2$, is

a) n if n is even b) 1 if n is odd c) 0 for any $n\in N$ d) None of these

111. Consider the following statements:

1. If x be real, then $-\frac{9}{2}\leq\frac{3x-4}{x^2+1}\leq\frac{1}{2}$

2. If x is real, then the greatest value of $\frac{x^2+14x+9}{x^2+2x+3}$ is 4

3. If $ax^2+bx+c=0; a\neq 0, a,b,c\in R$ has no real roots, then $(a+b+c)c>0$

Which of these is/ are correct?

a) Only (1) b) Only(2) c) Only (3) d) All of these

112. If r is a real number such that $|r|<1$ and if $a=5(1-r)$, then

a) $0<a<5$ b) $-5<a<5$ c) $0<a<10$ d) $0\leq a<10$

113. The number of integral roots of the equation $e^{x-8}+2x-17=0$, is

a) 1 b) 2 c) 4 d) 8

114. The product of real roots of the equation $x^2+18x+30=2\sqrt{x^2+18x+45}$, is

a) 720 b) 20 c) 36 d) None of these

115. The set of values of x satisfying $2\leq|x-3|<4$ is

a) $(-1,1]\cup[5,7)$ b) $-4\leq x\leq 2$

c) $-1<x<7$ or $x\geq 5$ d) $x<7$ or $x\geq 5$

116. Let $x=\left[\frac{a+2b}{a+b}\right]$ and $y=\frac{a}{b}$, where a and b are positive integers. If $y^2>2$, then

a) $x^2\leq 2$ b) $x^2<2$ c) $x^2>2$ d) $x^2\geq 2$

117. The least value of $5^{\sin x-1}+5^{-\sin x-1}$ is

a) 10 b) $\frac{5}{2}$ c) $\frac{2}{5}$ d) $\frac{1}{5}$

118. If $x^2+2x+n>10$ for all real numbers x, then which of the following conditions is true?

a) $n<11$ b) $n=10$ c) $n=11$ d) $n>11$

119. The minimum value of $P=bcx+cay+abz$, when $xyz=abc$, is

a) $3abc$ b) $6abc$ c) abc d) $4abc$

120. If $a_i>0$ for $i=1,2,\ldots,n$ and $a_1a_2\ldots a_n=1$, then minimum value of $(1+a_1)(1+a_2)\ldots(1+a_n)$ is

a) $2^{n/2}$ b) 2^n c) 2^{2n} d) 1

121. If a,b,c are positive real numbers such that $a+b+c=2$ then, which one of the following is true?

a) $(2-a)(2-b)(2-c)\geq 8abc$

b) $\frac{1}{a}+\frac{1}{b}+\frac{1}{c}\geq 2$

c) $(2-a)(2-b)(2-c)<8abc$

d) $\frac{1}{a}+\frac{1}{b}+\frac{1}{c}=2$

122. If x,y,z are positive real numbers such that $x^2+y^2+z^2=27$, then $x^3+y^3+z^3$ has

a) Minimum value 81 b) Maximum value 81 c) Minimum value 27 d) Maximum value 27

123. If x satisfies the inequations $2x-7<11, 3x+4<-5$, then x lies in the interval

a) $(-\infty,3)$ b) $(-\infty,2)$ c) $(-\infty,-3)$ d) $(-\infty,\infty)$

124. $x^8-x^5-\frac{1}{x}+\frac{1}{x^4}>0$, is satisfied for

a) Only positive values of x

b) Only negative values of x

c) All real numbers except zero

d) Only for $x>1$

125. The solution set of the inequation $5^{(1/4)(\log_5 x)^2}\geq 5\,x^{(1/5)(\log_5 x)}$, is

a) $(0,5^{-2\sqrt{5}}]$ b) $[5^{2\sqrt{5}},\infty)$ c) $(0,5^{-2\sqrt{5}}]\cup[5^{2\sqrt{5}},\infty)$ d) $(0,\infty)$

126. If $a+b=8$, then ab is greater when

a) $a=4, b=4$ b) $a=3, b=5$ c) $a=6, b=2$ d) None of these

127. The number of solutions of the equation $\cos x + |x| = 0$ is

a) 0 b) 1 c) 2 d) 3

128. If $0 < x < \frac{\pi}{2}$, then the minimum value of $\frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x}$ is

a) $\sqrt{3}$ b) $\frac{1}{2}$ c) $\frac{1}{3}$ d) 1

129. If $x^2 + 4ax + 2 > 0$ for all values of x, then a lies in the interval

a) (-2, 4) b) (1, 2) c) $(-\sqrt{2}, \sqrt{2})$ d) $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

130. If a and b are two different positive real numbers then which of the following statement is true?

a) $2\sqrt{ab} > a+b$ b) $2\sqrt{ab} < a+b$ c) $2\sqrt{ab} = a+b$ d) None of these

131. The number of negative integral solutions of $x^2 \cdot 2^{x+1} + 2^{|x-3|+2} = x^2 \cdot 2^{|x-3|+4} + 2^{x-1}$, is

a) None b) Only one c) Two d) Four

132. The number of ordered 4-tuples (x, y, z, w) where $x, y, z, w \in [0,10]$ which satisfy the inequality $2^{\sin^2 x} \times 3^{\cos^2 y} \times 4^{\sin^2 z} \times 5^{\cos^2 w} \geq 120$, is

a) 81 b) 144 c) 0 d) Infinite

133. If $a > 1, b > 1, c > 1, d > 1$, then the minimum value of $\log_b a + \log_a b + \log_d c + \log_c d$ is

a) 1 b) 2 c) 3 d) 4

134. The solution set of inequation$\log_{1/3}(2^{x+2} - 4^x) \geq -2$, is

a) $(-\infty, 2-\sqrt{13})$ b) $(-\infty, 2+\sqrt{13})$ c) $(-\infty, 2)$ d) None of these

135. If $\frac{2x}{2x^2+5x+2} > \frac{1}{x+1}$, then

a) $-2 > x > -1$ b) $-2 \geq x \geq -1$ c) $-2 < x < -1$ d) $-2 < x \leq -1$

136. The number of real solutions of $\log_2 x + |x| = 0$, is

a) 0 b) 1 c) 3 d) None of these

137. If $xyz = abc$, then the least value of $bcx + cay + abz$ is

a) $3abc$ b) $6abc$ c) abc d) $4abc$

138. The number of solution(s) of the inequation$\sqrt{3x^2+6x+7} + \sqrt{5x^2+10x+14} \leq 4 - 2x - x^2$, is

a) 1 b) 2 c) 4 d) Infinitely many

139. A stick of length 20 units is to be divided into n parts so that the product of the lengths of the parts is greater than unity. The maximum possible value of n is

a) 18 b) 19 c) 20 d) 21

140. If a, b, c are different positive real number such that $(b+c-a), (c+a-b)$and $(a+b-c)$ are positive, then $(a+b-c)(b+c-a)(c+a-b) - abc$ is

a) Positive b) Negative c) Non-positive d) Non-negative

141. $\log_{16} x^3 + \left(\log_2 \sqrt{x}\right)^2 < 1$ iff x lies in

a) (2, 16) b) (0,1/16) c) (1/16 ,2) d) None of these

142. If $\log_{\cos x} \sin x > 2$ and $0 < x < 3\pi$,then $\sin x$lies in the interval

a) $\left[\frac{\sqrt{5}-1}{2}, 1\right]$ b) $\left[0, \frac{\sqrt{5}-1}{2}\right]$ c) $\left[0, \frac{1}{2}\right]$ d) None of these

143. If $f(x) = x^2 + 2bx + 2c^2$ and $g(x) = -x^2 - 2cx + b^2$ such that $\min f(x) > \max g(x)$, then the relation between b and c, is

a) No real value of b and c b) $0 < c < b\sqrt{2}$ c) $|c| < |b|\sqrt{2}$ d) $|c| > |b|\sqrt{2}$

144. If the sum of the greatest integer less than or equal to x and the least integer greater than or equal to x is 5, then the solution set for x is

a) (2, 3) b) (0, 5) c) [5, 6) d) None of these

145. The total number of roots of the equation $|x - x^2 - 1| = |2x - 3 - x^2|$ is

a) 1 b) 2 c) 0 d) Infinitely many

146. For $\frac{|x-1|}{x+2} < 1, x$ lies in the interval

a) $(-\infty,-2) \cup \left(-\frac{1}{2},\infty\right)$ b) $(-\infty,1) \cup [2,3]$ c) $(-\infty,-4)$ d) $\left[-\frac{1}{2},1\right]$

147. Number of integer solutions of $\frac{x+2}{x^2+1} > \frac{1}{2}$ is

a) 0 b) 1 c) 2 d) 3

148. Solution of the inequality $\tan\left(x+\frac{\pi}{3}\right) \geq 1$ is

a) $\left(n\pi+\frac{\pi}{12}, n\pi+\frac{\pi}{6}\right)$ b) $\left(n\pi-\frac{\pi}{12}, n\pi+\frac{\pi}{6}\right)$ c) $\left(n\pi-\frac{\pi}{6}, n\pi-\frac{\pi}{12}\right)$ d) None of these

149. If $0 < a < 1$, then the solution set of the inequation $\frac{1+(\log_a x)^2}{1+(\log_a x)} > 1$, is

a) $(1, 1/a)$ b) $(0, a)$ c) $(1, 1/a) \cup (0, a)$ d) None of these

150. Let $x = \frac{a+2b}{a+b}$ and $y = \frac{a}{b}$, wherer a and b are positive integers. If $y^2 > 2$, then

a) $x^2 \leq 2$ b) $x^2 < 2$ c) $x^2 > 2$ d) $x^2 \geq 2$

151. The minimum value of $|\sin x + \cos x + \tan x + \sec x + \operatorname{cosec} x + \cot x|$ is

a) $2\sqrt{2}-1$ b) $2\sqrt{2}+1$ c) $\sqrt{2}-1$ d) $\sqrt{2}+1$

152. If for $x \in R, \frac{1}{3} < \frac{x^2-2x+4}{x^2+2x+4} < 3$, then $\frac{9\cdot 3^{2x}-6\cdot 3^x+4}{9\cdot 3^{2x}+6\cdot 3^x+4}$ lies between

a) $\frac{1}{2}$ and 2 b) $\frac{1}{3}$ and 3 c) 0 and 2 d) None of these

153. The minimum value of $4^x + 4^{1-x}, x \in R$, is

a) 1 b) 2 c) 4 d) None of these

154. The number of real solutions of the equation $3^{-|x|} - 2^{|x|} = 0$, is

a) 0 b) 1 c) 2 d) None of these

155. The number of real roots of the equation $1 + 3^{x/2} = 2^x$, is

a) 0 b) 1 c) 2 d) None of these

156. If n is even and $n \geq 4, x_1, x_2, \ldots, x_n \geq 0$ and $x_1 + x_2 + \cdots + x_n = 1$, then $P = x_1x_2 + x_2x_3 + \cdots + x_n - x_n$ cannot exceed

a) $\frac{1}{n+1}$ b) $\frac{1}{n+2}$ c) $\frac{1}{2n}$ d) None of these

157. The number of real solutions of the equation $e^{-x} = x$, is

a) 0 b) 1 c) 2 d) None of these

158. The solution set contained in R of the inequation $3^x + 3^{1-x} - 4 < 0$, is

a) $(1,3)$ b) $(0,1)$ c) $(1,2)$ d) $(0,2)$

159. The solution of the inequation $2x^2 + 3x - 9 \leq 0$ is given by

a) $\frac{3}{2} \leq x \leq 3$ b) $-3 \leq x \leq \frac{3}{2}$ c) $-3 \leq x \leq 3$ d) $\frac{3}{2} \leq x \leq 2$

160. If $0 < \theta < \pi$, then the minimum value of $\sin^5\theta + \operatorname{cosec}^5\theta$ is

a) 0 b) 1 c) 2 d) None of these

1 **(a)**

We have,

$3^x + 2^{2x} \geq 5^x$

$\Rightarrow \left(\frac{3}{5}\right)^x + \left(\frac{4}{5}\right)^x \geq 1$

$\Rightarrow \left(\frac{3}{5}\right)^x + \left(\frac{4}{5}\right)^x \geq \left(\frac{3}{5}\right)^2 + \left(\frac{4}{5}\right)^2$

$\Rightarrow x \leq 2 \Rightarrow x$

$\in (-\infty, 2]$ [If $a^x + b^x \geq 1$ and $a^2 + b^2 = 1$, then $x \in (-\infty, 2)$]

2 **(c)**

$x^2 - 3|x| + 2 < 0$

$\Rightarrow |x|^2 - 3|x| + 2 < 0$

$\Rightarrow (|x| - 1)(|x| - 2) < 0$

$\Rightarrow 1 < |x| < 2$

$\Rightarrow -2 < x < -1$ or $1 < x < 2$

$\therefore x \in (-2, -1) \cup (1,2)$

3 **(d)**

We have, $2^x + 2^{|x|} \geq 2\sqrt{2}$

If $x \geq 0$, then $2^x + 2^x \geq 2\sqrt{2}$

$\Rightarrow 2^x \geq \sqrt{2} \Rightarrow x \geq \frac{1}{2}$

and if $x < 0$, then $2^x + 2^{-x} \geq 2\sqrt{2}$

$\Rightarrow t + \frac{1}{t} \geq 2\sqrt{2}$ (where $t = 2^x$)

$\Rightarrow t^2 - 2\sqrt{2}t + 1 \geq 0$

$\Rightarrow \left(t - \left(\sqrt{2} - 1\right)\right)\left(t - \left(\sqrt{2} + 1\right)\right) \geq 0$

$\Rightarrow t \leq \sqrt{2} - 1$ or $t \geq \sqrt{2} + 1$

But $t > 0$

$\Rightarrow 0 < 2^x \leq \sqrt{2} - 1$

Or $2^x \geq \sqrt{2} + 1$

$\Rightarrow -\infty < x \leq \log_2(\sqrt{2} - 1)$

Or $x \geq \log_2\left(\sqrt{2} + 1\right)$

Which is not possible, because $x > 0$

$\therefore x \in (-\infty, \log_2(\sqrt{2} - 1)) \cup \left[\frac{1}{2}, \infty\right)$

4 **(b)**

Using G. M. $\leq$ A. M., we have

$\sin x_i \cos x_{i+1} \leq \frac{\sin^2 x_i + \cos^2 x_{i+1}}{2}$ for $i = 1,2,3, \ldots, n$,

where $x_{n+1} = x_1$

$\therefore \sin x_1 \cos x_2 + \sin x_2 \cos x_3 + \cdots + \sin x_n \cos x_{n+1}$

$\leq \frac{\sin^2 x_1 + \cos^2 x_2}{2} + \frac{\sin^2 x_2 + \cos^2 x_3}{2} + \cdots + \frac{\sin^2 x_n + \cos^2 x_1}{2}$

$\Rightarrow \sin x_1 \cos x_2 + \sin x_2 \cos x_3 + \cdots + \sin x_n \cos x_1 \leq \frac{n}{2}$

5 **(c)**

We have, $x^2 + (a + b)x + ab < 0$

$\Rightarrow (x + a)(x + b) < 0$

$\Rightarrow -b < x < -a$

6 **(a)**

Given, $\log_{10}(x^3 + y^3) - \log_{10}(x^2 + y^2 - xy) \leq 2$

$\Rightarrow \log_{10} \frac{(x^3 + y^3)}{x^2 + y^2 - xy} \leq 2$

$\Rightarrow \log_{10}(x + y) \leq 2 \Rightarrow x + y \leq 100$

Using AM $\geq$ GM

$\therefore \frac{x + y}{2} \geq \sqrt{xy}$

$\Rightarrow \sqrt{xy} \leq \frac{x + y}{2} \leq \frac{100}{2}$

$\Rightarrow xy \leq 2500$

7 **(c)**

By trial,

$3^{x/2} + 2^x \leq 25$ for $x = 1,2,3,4$

But $3^{x/2} + 2^x > 25$ for $x > 4$

Hence, solution set for $3^{x/2} + 2^x > 25$ is $(4, \infty)$

8 **(c)**

Since,

AM $\geq$ GM

$\Rightarrow \frac{a + b}{2} \geq \sqrt{ab}$

$\Rightarrow \frac{a+b}{2} \geq \sqrt{4}$ ($\because ab = 4$, given)

$\Rightarrow a + b \geq 4$

9 **(a)**

When $x > 0, P_n(x) > 0$ and so $P_n(x) = 0$ can have no positive real roots

Now,

$P_n(x) = 1 + 2x + 3x^2 + \cdots + (n+1)x^n$

$\Rightarrow xP_n(x) = x + 2x^2 + 3x^3 + \cdots + nx^n + (n+1)x^{n+1}$

$\Rightarrow (1-x)P_n(x) = 1 + x + x^2 + \cdots + x^n - (n+1)x^{n+1}$

$\Rightarrow P_n(x) = \dfrac{1-(n+2)x^{n+1}+(n+1)x^{n+2}}{(1-x)^2}$

For negative values of $x, P_n(x)$ will vanish whenever

$f(x) = 1-(n+2)x^{n+1}+(n+1)x^{n+2} = 0$

Now,

$f(-x) = 1-(n+2)(-1)^{n+1}x^{n+1} + (n+1)(-1)^{n+2}x^{n+2}$ ...(i)

If n is even, there is no change of sign in this expression and so there is no negative real root of $f(x)$

10 **(c)**

$(x-1)(x^2-5x+7) < (x-1)$

$\Rightarrow (x-1)(x^2-5x+6) < 0$

$\Rightarrow (x-1)(x-2)(x-3) < 0$

$\therefore x \in (-\infty, 1) \cup (2, 3)$

11 **(b)**

Using A. M. $\geq$ G. M., we have

$\dfrac{x}{n} \geq \{\log_{2^2} 2 \times \log_{2^3} 2^2 \times \log_{2^4} 2^3 \times \ldots \times \log_{2^{n+1}} 2^n\}^{1/n}$

$\Rightarrow x \geq n(\log_{2^{n+1}} 2)^{1/n}$

$\Rightarrow x \geq n\left(\dfrac{1}{n+1}\log_2 2\right)^{1/n} \Rightarrow x \geq n\left(\dfrac{1}{n+1}\right)^{1/n}$

12 **(c)**

We have,

$\frac{1}{2}\left|a + \frac{11}{a}\right| \geq \sqrt{11}$, equality holding iff $a = \pm\sqrt{11}$

$\therefore |x| \geq \sqrt{11}, |y| \geq \sqrt{11}, |z| \geq \sqrt{11}, |t| \geq \sqrt{11}$

Let $x \geq 0$, then $x \geq \sqrt{11}, y \geq \sqrt{11}, z \geq \sqrt{11}$ and $t \geq \sqrt{11}$

Now, $y - \sqrt{11} = \dfrac{1}{2}\left(\dfrac{11}{x} + x\right) - \sqrt{11}$

$\Rightarrow y - \sqrt{11} = \dfrac{\left(x-\sqrt{11}\right)^2}{2x} = \left(\dfrac{x-\sqrt{11}}{2x}\right)(x-\sqrt{11})$

$\Rightarrow y - \sqrt{11} = \dfrac{1}{2}\left(1 - \dfrac{\sqrt{11}}{x}\right)(x-\sqrt{11}) < (x-\sqrt{11})$

$\Rightarrow y < x$ i. e. $x > y$

Similarly, we have

$y > z, z > t$ and $t > x \Rightarrow y > x$

$\therefore x = y = z = t = \sqrt{11}$ is the only solution for $x > 0$

We observe that (x, y, z, t) is a solution iff $(-x, -y, -z, -t)$ is a solution

Thus, $x = y = z = t = -\sqrt{11}$ is the only other solution

13 **(b)**

Given, $3 + \frac{3}{3^x} - 4 < 0 \Rightarrow 3^{2x} + 3 - 4.3^x < 0$

$\Rightarrow (3^x - 1)(3^x - 3) < 0$

$1 < 3^x < 3 \Rightarrow 0 < x < 1$

$\therefore$ The solution set is $(0, 1)$

14 **(b)**

We have, $|a| \leq 1$ and $a + b = 1$

i. e. $-1 \leq a \leq 1$ and $b = 1 - a$

$\Rightarrow -1 \leq a \leq 1$ and $0 \leq b \leq 2 \Rightarrow -2 \leq ab \leq 2$ ...(i)

Now,

$ab \leq \left(\frac{a+b}{2}\right)^2 \Rightarrow ab \leq \frac{1}{4}$ ...(ii)

From (i) and (ii), we have

$-2 \leq ab \leq \dfrac{1}{4} \Rightarrow ab \in \left[-\dfrac{2,1}{4}\right]$

15 **(d)**

Given, $\sqrt{(3x+1)^2} < (2-x)$

$\Rightarrow (3x+1) < 2 - x$

$\Rightarrow 3x + 1 < 2 - x \Rightarrow x < \dfrac{1}{4}$

16 **(c)**

Given, inequality can be rewritten as $\left(\frac{5}{13}\right)^x + \left(\frac{12}{13}\right)^x \geq 1$

$\therefore \cos^x \alpha + \sin^x \alpha \geq 1$

Where, $\cos \alpha = \frac{5}{13}$

If $x = 2$, the above equality holds

If $x < 2$ both $\cos\alpha$ and $\sin\alpha$ increases in positive fraction

Hence, above inequality holds for $x \in (-\infty, 2]$

17 **(b)**

Let $f(x) = \log_e \frac{x-2}{x-3}$

$f(x)$ is defined either $(x-2)>0, (x-3)>0$ or $(x-2)<0$
$(x-3)<0$ or $x \neq 2,3$
$\Rightarrow f(x)$ is defined either $x>3$ or $x<2$ or $x \neq 2,3$
$ie, x \in (-\infty, 2) \cup (3, \infty)$

18 **(c)**
$3 \leq 3t-18 \leq 18$
$\Rightarrow \ 21 \leq 3t \leq 36$
$\Rightarrow 7 \leq t \leq 12$
$\Rightarrow 8 \leq t+1 \leq 13$

19 **(b)**
$\because f(-1)<1 \Rightarrow a-b+c<1$ ...(i)

and $f(1)>-1, f(3)<-4$, then

$a+b+c>-1$ ...(ii)

$9a+3b+c<-4$ ...(iii)

From Eq. (ii),

$-a-b-c<1$ ....(iv)

On solving Eqs. (i), (iii) and (iv), we get $a<-\frac{1}{8} \Rightarrow$ a is negative

20 **(d)**
Given, $\frac{2x+3}{2x-9}<0$
$\Rightarrow 2x+3<0$ and $2x-9>0$
Or $2x+3>0$ and $2x-9<0$ and $x \neq \frac{9}{2}$
$\Rightarrow \ x<-\frac{3}{2}$ and $x>\frac{9}{2}$ or $x>-\frac{3}{2}$ and $x<\frac{9}{2}$ and $x \neq \frac{9}{2}$
$\Rightarrow x \in \left(-\frac{3}{2}, \frac{9}{2}\right)$

21 **(b)**
We have,
$\sqrt{x^2+\sqrt{x^2+11}}-\sqrt{x^2-\sqrt{x^2+11}}=4$
Putting $x^2+11=t^2$, we get
$\sqrt{t^2+t-11}+\sqrt{t^2-t-11}=4$ ...(i)
But, $(t^2+t-11)-(t^2-t-11)=2t$ ...(ii)
Dividing (ii) by (i), we get
$\sqrt{t^2+t-11}-\sqrt{t^2-t-11}=\frac{t}{2}$ ...(iii)
Adding (i) and (iii), we get
$2\sqrt{t^2+t-11}=4+\frac{t}{2}$
$\Rightarrow t^2+t-11=4+t+\frac{t^2}{16}$
$\Rightarrow t^2=16 \Rightarrow t=4 \left[\because t=\sqrt{x^2+11}>0\right]$
$\therefore x=\pm\sqrt{5}$

22 **(a)**
We have,
$\log_{x-3}(x^3-3x^2-4x+8)=3$ ...(i)
$\Rightarrow x^3-3x^2-4x+8=(x-3)^3$
$\Rightarrow 6x^2-31x+35=0 \Rightarrow (3x-5)(2x-7)=0$
$\Rightarrow x=\frac{5}{3}, \frac{7}{2}$
The equation (i) exists, if
$x-3>0, x-3 \neq 1$ and $x^3-3x^2-4x+8>0$
Clearly, $x=\frac{7}{2}$ satisfies these conditions

23 **(a)**
Curves $y=\log_{0.5} x$ and $y=|x|$ intersect at one point in first quadrant. So, the equation $\log_{0.5} x=|x|$ has one real root

25 **(b)**
$\cos^x \alpha+\sin^x \alpha \geq 1$

Equality holds when $x=2$

If $x<2$, both $\cos\alpha$ and $\sin\alpha$ are increasing

$\therefore \cos^x \alpha+\sin^x \alpha>1$, if $x<2$

If $x>2$, then $\cos^x \alpha+\sin^x \alpha<1$

$\therefore x \in (-\infty, 2]$

27 **(a)**
We have,
$2\cos(e^x)=3^x+3^{-x}$
We observe that $2\cos(e^x)<2$ and $3^x+3^{-x} \geq 2$.
So, the given equation has no solution

28 **(b)**
Graphs of $y=1-x$ and $y=[\cos x]$ cut each other at point $(0,1)$ and at a point whose x-coordinate lie in $(\pi/2, \pi)$. So, the given equation has two real roots

30 **(b)**
If $a^2+b^2=1$, then $a^x+b^x \geq 1$ is true for all $x \in (-\infty, 2]$
$\therefore (\sin\alpha)^x+(\cos\alpha)^x \geq 1 \Rightarrow x \in (-\infty, 2]$

31 **(a)**
If $f(x)>0$,then $D<0$
$4a^2-4(10-3a)<0$
$\Rightarrow \ (a+5)(a-2)<0$
$\Rightarrow \ -5<a<2$

32 **(b)**
The given inequality is

$49.4-\left(\frac{27-x}{10}\right)<47.4-\left(\frac{27-9x}{10}\right)$

$\Rightarrow 49.4 - 47.4 < \left(\frac{27-x}{10}\right) - \left(\frac{27-9x}{10}\right)$

$\Rightarrow 2 < \frac{8x}{10} \Rightarrow x > \frac{5}{2}$

$\therefore$ Least integer is 3

33 **(a)**

Since, AM $\geq$ GM

$\therefore \frac{a^2+b^2}{2} \geq \sqrt{a^2b^2} = ab, \frac{b^2+c^2}{2} \geq bc$

and $\frac{c^2+a^2}{2} \geq ca$

On adding, we get

$a^2 + b^2 + c^2 \geq ab + bc + ca$

$\Rightarrow (a)$holds

Next, $\frac{b+c}{2} \geq \sqrt{bc}$, $\frac{c+a}{2} \geq \sqrt{ca}, \frac{a+b}{2} \geq \sqrt{ab}$

$\Rightarrow \left(\frac{b+c}{2}\right)\left(\frac{c+a}{2}\right)\left(\frac{a+b}{2}\right) \geq \sqrt{a^2b^2c^2}$

$\Rightarrow (b+c)(c+a)(a+b) \geq 8abc$

$\Rightarrow (b)$ does not hold

Again, $\frac{1}{3}\left(\frac{a}{b} + \frac{b}{c} + \frac{c}{a}\right) \geq \left(\frac{a}{b} \cdot \frac{b}{c} \cdot \frac{c}{a}\right)^{1/3}$

$\Rightarrow \frac{a}{b} + \frac{b}{c} + \frac{c}{a} \geq 3$

$\Rightarrow (c)$does not hold

Again, $\frac{a^3+b^3+c^3}{3} \geq (a^3b^3c^3)^{1/3}$

$\Rightarrow a^3 + b^3 + c^3 \geq 3\,abc$

$\Rightarrow (d)$ does not hold

34 **(d)**

If s is the semi-perimeter of a cyclic quadrilateral of sides a, b, c and d units in length, then its area A is given by

$A = \sqrt{(s-a)(s-b)(s-c)(s-d)}$

Using A. M. $\geq$ G. M., we have

$\frac{s-a+s-b+s-c+s-d}{4} \geq \{(s-a)(s-b)(s-c)(s-d)\}^{1/4}$

$\Rightarrow \frac{4s-2s}{4} \geq \sqrt{A} \Rightarrow 2s \geq 4\sqrt{A}$

Hence, the least perimeter is $4\sqrt{A}$

35 **(c)**

Two curves

$y = [\sin x + \cos x]$

and, $y = 3 + [-\sin x] + [-\cos x]$

$= 1 + [\sin x] + [\cos x]$

intersect at infinitely many points in $[0, 2\pi]$

So, the given equation has infinitely many solutions

36 **(b)**

Two curves $y = 3^{|x|}$ and $y = \left|2 - |x|\right|$ intersect at two points only. So, the equation $3^{|x|} = \left|2 - |x|\right|$ has only two real roots

37 **(a)**

Since, angle C is obtuse, angle A and B are actute

$\therefore \tan C < 0$and $\tan A > 0, \tan B > 0$

Now, $\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A\ \tan B}$

$\Rightarrow \tan(\pi - C) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$

$\Rightarrow -\tan C = \frac{\tan A + \tan B}{1 - \tan A \tan B}$

$\Rightarrow 1 - \tan A \tan B > 0$ ($\because$Numerator are positive)

$\Rightarrow \tan A \tan B < 1$

38 **(a)**

We have,

$x + y + z = 4$ and $x^2 + y^2 + z^2 = 6$

$\Rightarrow y + z = 4 - x$ and $y^2 + z^2 = 6 - x^2$

$\therefore yz = \frac{1}{2}\{(y+z)^2 - (y^2+z^2)\}$

$= \frac{1}{2}\{(4-x)^2 - (6-x)^2\}$

$\Rightarrow yz = x^2 - 4x + 5$

Thus, y and z are roots of the equation

$t^2 - (4-x)\,t + x^2 - 4x + 5 = 0$

As y, z are real

$\therefore (4-x)^2 - 4(x^2 - 4x + 5) \geq 0$

$\Rightarrow 3x^2 - 8x + 4 \leq 0 \Rightarrow \frac{2}{3} \leq x \leq 2$

40 **(b)**

We have,

$3^{x/2} + 2^x > 25 \Rightarrow 3^{x/2} + 4^{x/2} > 25$

Clearly, $x \in (4, \infty)$ satisfies the above inequation

41 **(c)**

We have,

$27^{1/x} + 12^{1/x} = 2 \times 8^{1/x}$

$\Rightarrow 3^{3/x} + 2^{2/x} \times 3^{1/x} = 2 \times 2^{3/x}$

$\Rightarrow \left(\frac{3}{2}\right)^{3/x} + \left(\frac{3}{2}\right)^{1/x} = 2$

$\Rightarrow y^3 + y - 2 = 0$, where $y = \left(\frac{3}{2}\right)^{1/x}$

$\Rightarrow (y-1)(y^2+y-2) = 0$

$\Rightarrow y = 1, y = -2 \Rightarrow \left(\frac{3}{2}\right)^{1/x} = 1 \Rightarrow \left(\frac{3}{2}\right)^{1/x} = \left(\frac{3}{2}\right)^0$

But, there is no value of x for which $\frac{1}{x}$ is zero

Hence, the given equation has no solution

42 **(c)**

Let x_1, x_2, x_3 and x_4 be four positive roots of the equation $x^4 - 8x^3 + bx^2 + cx + 16 = 0$. Then,

$x_1 + x_2 + x_3 + x_4 = 8$ and $x_1x_2x_3x_4 = 16$

$\Rightarrow \frac{x_1 + x_2 + x_3 + x_4}{4} = 2$ and $(x_1x_2x_3x_4)^{1/4} = 2$

$\Rightarrow$ A. M. and G. M. of x_1, x_2, x_3 and x_4 are equal

$\Rightarrow x_1 = x_2 = x_3 = x_4$

$\Rightarrow x_1 = x_2 = x_3 = x_4 = 2$

$\therefore x^4 - 8x^3 + bx^2 + cx + 16 = (x-2)^4$

$\Rightarrow b = {}^4C_2 \times 2^2 = 24$ and $c = -{}^4C_3 \times 2^3 = -32$

43 **(a)**

We have, $3 < |x| < 6 \Rightarrow -6 < x < -3$ or $3 < x < 6$

$\therefore x \in (-6, -3) \cup (3, 6)$

44 **(a)**

We have,

$a^4 + b^4 - a^3 b - ab^3 = a^3(a-b) - b^3(a-b)$

$= (a^3 - b^3)(a-b)$

$\Rightarrow a^4 + b^4 - a^3b - ab^3 > 0$ $\left[\because a^3 - b^3 \text{ and } a - b \text{ are of the same sign}\right]$

$\Rightarrow a^4 + b^4 > a^3 b + ab^3$

45 **(a)**

The given inequation is

$4^{-x+0.5} - 7 \cdot 2^{-x} < 4, x \in R$

Let $2^{-x} = t$

$\therefore 2t^2 - 7t < 4$

$\Rightarrow 2t^2 - 7t - 4 < 0$

$\Rightarrow (2t+1)(t-4) < 0$

$\Rightarrow -\frac{1}{2} < t < 4$

$\Rightarrow 0 < t < 4 \quad (\because t = 2^{-x} > 0)$

$\Rightarrow 0 < 2^{-x} < 2^2$

As 2^x is an increasing function – $x < 2$ or $x > -2$

$\therefore x = (-2, \infty)$

46 **(c)**

Given condition are $\frac{a}{b} > 1$ and $\frac{a}{c} < 0$

1. $a > 0$ iff $c < 0$ and also $b > 0$
2. $a < 0$ iff $c > 0$ and also $b < 0$

47 **(b)**

Proceeding as in the solution of Q. no. 10, we have

$(a+b)(b+c)(c+a) \geq 8abc$

$\Rightarrow (p-a)(p-b)(p-c) \geq 8abc \quad [\because a + b + c = p]$

48 **(a)**

We have,

$\frac{\left(1 + e^{x^2}\right)\sqrt{1+x^2}}{\sqrt{1+x^4} - x^2} = 1 + \cos x$

$\Rightarrow \left(1 + e^{x^2}\right)\sqrt{1+x^2}\left(\sqrt{1+x^4} + x^2\right) = 1 + \cos x$

Clearly, LHS ≥ 2 and RHS ≤ 2. So, the equation exists when each side is equal to 2. This is for $x = 0$ only. Hence, it has only one solution

50 **(c)**

Let $f(x) = c_0 + c_1x + c_2x^2 + \cdots + c_nx^n$. Then,

$f(b) - f(a) = 1$

$\Rightarrow c_1(b-a) + c_2(b^2 - a^2) + \cdots + c_n(b^n - a^n) = 1$

$\Rightarrow (b-a)\{c_1 + c_2(b+a) + \cdots + c_n(b^{n-1} + b^{n-2}a + \cdots + a^{n-1})\} = 1$

$\Rightarrow (b-a)I = 1$, where I is an integer

$\Rightarrow b - a = \pm 1$

52 **(d)**

Using, AM > GM

$\therefore \frac{a+b+c}{3} > \sqrt[3]{abc}$

$\Rightarrow a + b + c > 3$ ...(i)

$[\because abc = 1$ given$]$

Also, GM > HM

$\sqrt[3]{abc} > \frac{3}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}$

$\Rightarrow (1)^{1/3} > \frac{3abc}{bc + ac + ab}$

$\Rightarrow ab + bc + ac > 3$ ...(ii)

$\therefore$ From Eqs. (i) and (ii), we get

$a + b + c + ab + bc + ac > 6$

53 **(d)**

We have,

$\sin(2^x)\cos(2^x) = \frac{2^x + 2^{-x}}{2}$

$\Rightarrow \sin(2^{x+1}) = 2^x + 2^{-x}$

Clearly, RHS ≥ 2 and LHS lies between -1 and 1.
So, the given equation has no solution

54 **(d)**

$x^{12} - x^9 + x^4 - x + 1 > 0$

When $0 < x < 1; x^4 > x^9$ and $1 > x$

$\therefore x^{12} + (x^4 - x^9) + (1 - x) > 0$

$\Rightarrow$ Positive for all x

Again, when $x > 1$:

$x^{12} - x^9 + x^4 - x + 1 > 0$

$\therefore$ Largest interval $(0, \infty)$, also the above inequality is true for $x < 0$

56 **(d)**

$\because$ AM $\geq$ GM

$$\Rightarrow \frac{\frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x}}{2} \geq \left(\frac{\cos^3 x}{\sin x} \cdot \frac{\sin^3 x}{\cos x}\right)^{1/2}$$

$$\Rightarrow \frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x} \geq 2 \sin x \cos x \geq 1$$

Hence, option (d) is correct

57 **(a)**

We have,

$\sqrt{3x^2 + 6x + 7} = \sqrt{3(x+1)^2 + 4} \geq 2($
$= 2$ when $x = -1)$

$\sqrt{5x^2 + 10x + 14} = \sqrt{5(x+1)^2 + 9} \geq 3($
$= 3$ when $x = -1)$

and,

$4 - 2x - x^2 = 5 - (x+1)^2 \leq 5 (= 5$ when x
$= -1)$

Thus, LHS ≥ 5 and RHS ≤ 5

So, the given equation is valid when each sides is equal to 5.

This happens only when $x = -1$

Hence, the given equation has only one solution

58 **(d)**

$\big||x| - 1\big| < |1 - x|$

Case I $x \geq 0$

$\therefore$ Inequality (i) becomes $|x - 1| < x - 1$ or $|1 - x| < 1 - x$ which is not satisfied by any x, because

$|a| \geq \forall a \in R$

Case II $-1 \leq x < 0$

$\therefore$ Inequality (i) becomes $|-1 - x| < 1 - x$ or $|x + 1| < 1 - x$

Or $x + 1 < 1 - x$ or $x < 0$

Thus, inequality (i) is satisfied for $-1 \leq x < 0$

Case III $x < -1$

Inequality (i) becomes $|-1 - x| < 1 - x \Rightarrow |1 + x| < 1 - x$

$\Rightarrow -(1 + x) < 1 - x \Rightarrow -2 < 0$, which is true

So, solution set is $(-\infty, 0)$

59 **(c)**

Minimum value of $f(x)$
Is attained at $x = 3$
$\therefore$ Minimum value of $f(x) = 7$

60 **(a)**

$$\frac{x + 11}{x - 3} > 0$$

$\Rightarrow \quad (x - 3)(x + 11) > 0$
$\Rightarrow \quad x < -11, x > 3$
$\Rightarrow \quad x \in (-\infty, -1) \cup (3, \infty)$

61 **(b)**

$$2x - 1 = |x + 7| = \begin{cases} x + 7, \text{if } x \geq -7 \\ -(x + 7), \text{if } x < -7 \end{cases}$$

$\therefore$ If $x \geq -7, 2x - 1 = x + 7 \Rightarrow x = 8$

If $x < -7, 2x - 1 = -(x + 7)$

$\Rightarrow 3x = -6$

$\Rightarrow x = -2$, which is not possible

62 **(c)**

Let $f(x) = x^4 - 4x - 1$. Then, the number of changes of signs in $f(x)$ is 1. Therefore, $f(x)$ can have at most one positive real root

We have,

$f(2) > 0$ and $f(0) < 1$

Therefore, $f(x)$ has one positive real root between 1 and 2

63 **(c)**

$\log_{\sin\left(\frac{\pi}{3}\right)}(x^2 - 3x + 2) \geq 2$

$\Rightarrow x^2 - 3x + 2 \leq \frac{3}{4}$ (If $\log_a b = c \Rightarrow b = c^a$)

$$\Rightarrow x^2 - 3x + \frac{5}{4} \leq 0$$

$\Rightarrow 4x^2 - 12x + 5 \leq 0$

$\Rightarrow (2x-5)(2x-1) \leq 0$

$\Rightarrow \frac{1}{2} \leq x \leq \frac{5}{2}$ ...(i)

Also, $x^2 - 3x + 2 > 0$

$\Rightarrow (x-1)(x-2) > 0$

$\Rightarrow x < 1$ or $x > 2$ ...(ii)

From relation (i) and (ii), we get

$x \in \left[\frac{1}{2}, 1\right) \cup \left(2, \frac{5}{2}\right]$

64 **(c)**

Since, $(a-b)^2 + (b-c)^2 + (c-a)^2 \geq 0$

$\Rightarrow \ 2(a^2 + b^2 + c^2) \geq 2(ab + bc + ca)$

$\Rightarrow \ \frac{a^2 + b^2 + c^2}{(ab + bc + ca)} \geq 1$

$\Rightarrow \frac{a^2 + b^2 + c^2}{ab + bc + ca} + 2 \geq 3$

Hence, option (c) is correct

65 **(b)**

Given, $(x-1)(x^2 - 5x + 7) < (x-1)$

$\Rightarrow \ (x-1)(x^2 - 5x + 6) < 0$

$\Rightarrow \ (x-1)(x-2)(x-3) < 0$

$\Rightarrow \ x \in (-\infty, 1) \cup (2, 3)$

66 **(d)**

Given inequalities are

$x^2 - 3x - 10 < 0$ and $10x - x^2 - 16 > 0$

$\Rightarrow \ (x+2)(x-5) < 0$ and $(x-2)(x-8) < 0$

$\Rightarrow \ x \in (-2, 5)$ and $x \in (2, 8)$

$\Rightarrow \ x \in (2, 5)$

67 **(a)**

$\log_{1/3}(x^2 + x + 1) < -1 = \log_{1/3}\left(\frac{1}{3}\right)^{-1}$

$\Rightarrow x^2 + x + 1 > \left(\frac{1}{3}\right)^{-1}$

($\because$ where $0 < a < 1$, then $\log_a x < \log_a y \Rightarrow x > y$)

$\Rightarrow x^2 + x - 2 > 0 \Rightarrow (x+2)(x-1) > 0$

$\Rightarrow x \in (-\infty, -2) \cup (1, \infty)$

68 **(a)**

Let (α, β) be a solution of the system for some a. Then, $(-\alpha, \beta)$ is also a solution. So, the system will have unique solution only if

$\alpha = -\alpha \Rightarrow \alpha = 0$

Putting $x = \alpha = 0$ and $y = \beta$ in $x^2 + y^2 = 1$, we get $\beta = \pm 1$

Putting $x = \alpha = 0$ and $y = \beta$ in $2^{|x|} + |x| = y + x^2 + a$, we get

$\beta + a = 1 \Rightarrow a = 1 - \beta$

$\therefore a = 0$ when $\beta = 1$ and $a = 2$ when $\beta = -1$

CASE I $When\ a = 0$

In this case, given equations become

$2^{|x|} + |x| = y + x^2$ and $x^2 + y^2 = 1$

Now, $x^2 + y^2 = 1 \Rightarrow |x| \leq 1$ and $|y| \leq 2$

$\therefore 2^{|x|} + |x| = y + x^2$ and $1 + x^2 \geq y + x^2$

$\Rightarrow 2^{|x|} + |x| \leq 1 + x^2$

$\Rightarrow 2^{|x|} + |x| \leq 1 + |x| \quad [\because x^2 \leq |x| \text{when } |x| < 1]$

$\Rightarrow x = 0$

Putting $x = 0$ in $2^{|x|} + |x| = y + x^2$, we get $y = 1$

Thus, for $a = 0$, the system has unique solution $(0, 1)$

CASE II $When\ a = 2$

In this case, the system of equation is

$2^{|x|} + |x| = y + x^2 + 2$ and $x^2 + y^2 = 1$

Clearly, $(0, -1)$, $(1, 0)$ and $(-1, 0)$ satisfy these equations.

So, the system does not have unique solution

69 **(c)**

We have,

$\log_{1/3}(x^2 + x + 1) + 1 > 0$

$\Rightarrow \log_{1/3}(x^2 + x + 1) > -1$

$\Rightarrow x^2 + x + 1 < \left(\frac{1}{3}\right)^{-1}$

$\Rightarrow x^2 + x + 1 < 3$

$\Rightarrow x^2 + x - 2 < 0 \Rightarrow (x+2)(x-1) < 0 \Rightarrow x \in (-2, 1)$

70 **(d)**

We have,

$x^{(\log_{10} x)^2 - 3(\log_{10} x) + 1} > 10^3$

$\Rightarrow (\log_{10} x)^2 - 3(\log_{10} x) + 1 > \log_x 10^3$

$\Rightarrow (\log_{10} x)^2 - 3(\log_{10} x) + 1 > \frac{3}{\log_{10} x}$

$\Rightarrow \frac{(\log_{10} x)^3 - 3(\log_{10} x)^2 + (\log_{10} x) - 3}{\log_{10} x} > 0$

$\Rightarrow \frac{\{(\log_{10} x)^2 + 1\}(\log_{10} x - 3)}{\log_{10} x} > 0$

$\Rightarrow \frac{(\log_{10} x - 3)}{(\log_{10} x - 0)} > 0$

$\Rightarrow \log_{10} x < 0$ or, $\log_{10} x > 3$

$\Rightarrow x < 1$ or, $x > 10^3$

$\Rightarrow x \in (0,1) \cup (10^3, \infty)$ [$\because \log_{10} x$ is defined for $x > 0$]

71 **(a)**

Given, $\frac{3-|x|}{4-|x|} \geq 0$

$\Rightarrow \ 3-|x| \leq 0$ and $4-|x|<0$
Or $3-|x| \geq 0$ and $4-|x|>0$
$\Rightarrow \ |x| \geq 3$ and $|x|>4$
Or $|x| \leq 3$ and $|x|<4$
$\Rightarrow \ |x|>4$ or $|x| \leq 3$
$\Rightarrow (-\infty,-4) \cup [-3,3] \cup (4,\infty)$

72 **(a)**

Now, $3(a^2+b^2+c^2)-(a+b+c)^2$

$=2(a^2+b^2+c^2-bc-ca-ab)$

$=(b-c)^2+(c-a)^2+(a-b)^2 \geq 0$

$\Rightarrow 3(a^2+b^2+c^2) \geq (a+b+c)^2>9$

$\Rightarrow a^2+b^2+c^2>3 \Rightarrow$ (a)holds

Now, $a^6+b^6 \geq 12a^2b^2-64$

If $a^6+b^6+64 \geq 12a^2b^2$

$ie, \ \ a^6+b^6+2^6 \geq 3 \cdot 2^2 \cdot a^2b^2$

$ie,$ if $\frac{a^6+b^6+2^6}{3} \geq (2^6a^6b^6)^{1/3} \quad (\because$ AM $\geq$ GM)

$\Rightarrow (b)$does not hold

Again, since AM $\geq$ HM

$\therefore \frac{a+b+c}{3} \geq \frac{3}{\frac{1}{a}+\frac{1}{b}+\frac{1}{c}}$

$\Rightarrow \frac{\alpha}{3} \geq \frac{3}{\frac{1}{a}+\frac{1}{b}+\frac{1}{c}}$

$\Rightarrow \frac{1}{a}+\frac{1}{b}+\frac{1}{c} \geq \frac{9}{\alpha}$

$\Rightarrow (c\ )$ does not hold

73 **(c)**

Using A. M. $\geq$ G. M., we have

$\frac{a_1+a_2+a_3}{3} \geq (a_1a_2a_3)^{1/3}$ and $\frac{1}{3}(\frac{1}{a_1}+\frac{1}{a_2}+\frac{1}{a_3}) \geq \left(\frac{1}{a_1a_2a_3}\right)^{1/3}$

$\Rightarrow (a_1+a_2+a_3)\left(\frac{1}{a_1}+\frac{1}{a_2}+\frac{1}{a_3}\right) \geq 9$

74 **(a)**

We have,

$x-\frac{1}{x}=B$ and $x^2+\frac{1}{x^2}=A$

$\therefore \left(x-\frac{1}{x}\right)^2=B^2$

$\Rightarrow A-2=B^2 \Rightarrow A=B^2+2 \Rightarrow \frac{A}{B}=B+\frac{2}{B}$

But, A. M. $\geq$ G. M.

$\Rightarrow B+\frac{2}{B} \geq 2\sqrt{B \times \frac{2}{B}} \Rightarrow B+\frac{2}{B} \geq 2\sqrt{2} \Rightarrow \frac{A}{B} \geq 2\sqrt{2}$

Hence, the minimum value of $\frac{A}{B}$ is $2\sqrt{2}$

75 **(c)**

As discussed in the above problem, if n is odd, there is one change of sign in (i). Therefore, $f(x)$ can have at most one negative real root. In this case, we have

$f(-1)=-2n-2<0, f(0)=1>0$

So, the negative real root lies between -1 and 0

76 **(c)**

Given, ${}^{n+1}C_{n-2}-{}^{n+1}C_{n-1} \leq 50$

$\Rightarrow \frac{(n-1)!}{3!\,(n-2)!}-\frac{(n+1)!}{2!\,(n-1)!} \leq 50$

$\Rightarrow \frac{(n+1)!}{3!}\left[\frac{1}{(n-2)!}-\frac{3}{(n-1)!}\right] \leq 50$

$\Rightarrow (n+1)!\left(\frac{n-1-3}{(n-1)!}\right) \leq 300$

$\Rightarrow (n+1)n(n-4) \leq 300$

For $n=8$, it satisfy to the above inequality

But $n=1$ it does not satisfy the above inequality

77 **(d)**

We have,

$f(\theta)=\sec^2\theta+\cos^2\theta=(\sec\theta-\cos\theta)^2+2 \geq 2$

$\Rightarrow f(\theta) \in [2,\infty)$

78 **(d)**

Let $a_1, a_2, \ldots, a_n$ be n positive integers such that

$a_1a_2 \ldots a_n=n^n$

Since, AM $\geq$ GM

$\therefore \frac{a_1+a_2+\ldots+a_n}{n} \geq (a_1a_2 \ldots a_n)^{1/n}$

$\Rightarrow \frac{a_1+a_2+\ldots+a_n}{n} \geq n$

$\Rightarrow a_1+a_2+\ldots+a_n \geq n^2$

79 **(a)**

$\log_2(x^2-3x+18)<4$

$\Rightarrow x^2-3x+18<16 \quad (\text{If} \log_a b<c \Rightarrow b<a^c)$

$\Rightarrow x^2-3x+2<0$

$\Rightarrow (x-1)(x-2)<0$

$\Rightarrow x \in (1,2)$

80 **(d)**

We have,

$[x]^2 = [x+6]$

$\Rightarrow [x]^2 = [x] + 6$

$\Rightarrow [x]^2 - [x] - 6 = 0$

$\Rightarrow ([x]-3)([x]+2) = 0$

$\Rightarrow [x] = 3, [x] = -2$

$\Rightarrow x \in [3,4)$ or $x \in [-2,-1) \Rightarrow x \in [-2,-1) \cup [3,4)$

81 **(b)**

Using AM $\geq$ GM

$\frac{\frac{a}{b+c}+\frac{b}{c+a}+\frac{c}{a+b}}{3} \geq \sqrt[3]{\frac{abc}{(a+b)(b+c)(c+a)}}$ ...(i)

Again, using AM$\geq$GM

$\frac{a+b}{2} \geq \sqrt{ab}, \frac{b+c}{2} \geq \sqrt{bc}, \frac{c+a}{2} \geq \sqrt{ca}$

$\Rightarrow (a+b)(b+c)(c+a) \geq 8abc$

$\Rightarrow \sqrt[3]{\frac{abc}{(a+b)(b+c)(c+a)}} \leq \frac{1}{2}$

$\therefore$ From Eq. (i)

$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}$

82 **(a)**

Two curves $y = e^{|x|}$ and $y = |x|$ does not intersect. So, the equation $e^{|x|} - |x| = 0$ has no solution

83 **(b)**

$|p-q| = \begin{cases} p-q, p \geq q \\ q-p, p < q \end{cases}$

$\therefore \min(p,q) = \begin{cases} p, p < q \\ q, q < p \end{cases}$

$\Rightarrow$ RHS $= \frac{1}{2}(p+q-|p-q|)$, if $p > q$

$\Rightarrow \frac{1}{2}(p+q-p+q) = q$

and LHS $\min(p,q) = q$

$\therefore \min(p,q) = \frac{1}{2}(p+q-|p-q|)$

84 **(a)**

The given equation is

$1 + |e^x - 1| = e^x(e^x - 2)$

$\Rightarrow |e^x - 1| + 2 = (e^x - 1)^2$

$\Rightarrow |e^x - 1|^2 - (e^x - 1) - 2 = 0$

$\Rightarrow (|e^x - 1| - 2)(|e^x - 1| + 1) = 0$

$\Rightarrow |e^x - 1| - 2 = 0 \quad [\because |e^x - 1| + 1 \neq 0]$

$\Rightarrow e^x - 1 = \pm 2 \Rightarrow e^x = 3, -1$

$\Rightarrow e^x = 3 \Rightarrow x = \log_e 3$ $[\because e^x > 0$ for all $x]$

85 **(c)**

We have,

$\log_{\sin\frac{2\pi}{3}}(x^2 - 3x + 2) \geq 2$

$\Rightarrow (x^2 - 3x + 2) \leq \left(\frac{\sqrt{3}}{2}\right)^2$ and $x^2 - 3x + 2 > 0$

$\Rightarrow 4x^2 - 12x + 5 \leq 0$ and $x^2 - 3x + 2 > 0$

$\Rightarrow (2x-1)(2x-5) \leq 0$ and $(x-1)(x-2) > 0$

$\Rightarrow \frac{1}{2} \leq x \leq \frac{5}{2}$ and $x < 1$ or $x > 2$

$\Rightarrow x \in [1/2, 1) \cup (2, 5/2]$

86 **(c)**

We have, $x^2 + 6x - 27 > 0$

$\Rightarrow (x+9)(x-3) > 0 \Rightarrow x < -9$ or $x > 3$

$\Rightarrow x \in (-\infty, -9) \cup (3, \infty)$....(i)

And $x^2 - 3x - 4 < 0$

$\Rightarrow (x-4)(x+1) < 0$

$\Rightarrow -1 < x < 4$ ....(i)

From relations (i) and (ii), we get

$3 < x < 4$

87 **(d)**

We have,

$\frac{1}{2}\{(x+y) + |x-y|\} = x$

$\Rightarrow \frac{1}{2}\{(x+y) + |x-y|\} = \frac{1}{2}\{(x+y) + (x-y)\}$

$\Rightarrow |x-y| = x - y \Rightarrow x \geq y$

88 **(c)**

Given, $\frac{4x-1}{3x+1} - 1 \geq 0$

$\Rightarrow \frac{x-2}{3x+1} \geq 0$

$\Rightarrow x - 2 \geq 0$ and $3x + 1 > 0$

Or $x - 2 \leq 0$ and $3x + 1 < 0$

$\Rightarrow x \geq 2$ and $x < -\frac{1}{3}$

Or $x \leq 2$ and $x > -\frac{1}{3}$

$\Rightarrow x \in \left(-\infty, -\frac{1}{3}\right) \cup [2, \infty)$

89 **(c)**

Given inequality holds only, if

$\sin^2 \alpha_i = 1$ or $\alpha_i = \pm\frac{\pi}{2}, \frac{3\pi}{2}$; $(i = 2,3, \ldots, n)$

$\Rightarrow$ Number of solutions $= 3 \times 3 \times 3 \times \ldots \times (n-1)$ times

$= 3^{n-1}$

90 **(b)**

We have, $e^x = x(x+1), x < 0$

Consider the curves $y = e^x$ and $y = x(x+1)$ for $x < 0$. Graphs of these two curve intersect at exactly one point. So, the equation $e^x = x(x+1)$ has exactly one real root

91 **(b)**

Draw graphs of $y = 1 - x + [x]$ and $y = \frac{1}{x} - \frac{1}{[x]}$

These two curves intersect it infinitely many points

92 **(d)**

We have, $2^x + 2^{|x|} \geq 2\sqrt{2}$

Following cases arise:

CASE I $When x \geq 0$

In this case, we have

$$2^x + 2^x \geq 2\sqrt{2} \Rightarrow 2^x \geq 2^{1/2} \Rightarrow x \geq \frac{1}{2} \Rightarrow x \in \left[\frac{1}{2}, \infty\right)$$

CASE II $When x < 0$

In this case, we have

$2^x + 2^{-x} \geq 2\sqrt{2}$

$\Rightarrow (2^x)^2 - 2\sqrt{2} \times 2^x + 1 \geq 0$

$\Rightarrow \left(2^x - \sqrt{2}\right)^2 - 1 \geq 0$

$\Rightarrow \left(2^x - \sqrt{2} - 1\right)\left(2^x - \sqrt{2} + 1\right) \geq 0$

$\Rightarrow 2^x \leq \sqrt{2} - 1$ or, $2^x \geq \sqrt{2} + 1$

$\Rightarrow x \leq \log_2\left(\sqrt{2} - 1\right)$ or, $x \geq \log_2\left(\sqrt{2} + 1\right)$

$\Rightarrow x \leq \log_2\left(\sqrt{2} - 1\right) \Rightarrow x \in \left(-\infty, \log_2\left(\sqrt{2} - 1\right)\right)$

Hence, $x \in \left(-\infty, \log_2\left(\sqrt{2} - 1\right)\right) \cup [1/2, \infty)$

93 **(c)**

$$\sin^4\frac{x}{3} + \cos^4\frac{x}{3} > \frac{1}{2} \Rightarrow 1 - \frac{1}{2}\sin^2\frac{2x}{3} > \frac{1}{2}$$

$$\Rightarrow \sin^2\frac{2x}{3} < 1 \Rightarrow \frac{2x}{3} \in \left(R - (2n+1)\frac{\pi}{2}\right)$$

$$\Rightarrow x \in R - \left(\frac{3n\pi}{2} + \frac{3\pi}{4}\right); n \in I$$

94 **(a)**

We have,

$(\log_5 x)^2 + (\log_5 x) < 2$

$\Rightarrow (\log_5 x)^2 + (\log_5 x) - 2 < 0$

$\Rightarrow (\log_5 x + 2)(\log_5 x - 1) < 0$

$$\Rightarrow -2 < \log_5 x < 1 \Rightarrow 5^{-1} < x < 5 \Rightarrow x \in \left(\frac{1}{25}, 5\right)$$

96 **(d)**

$\because \sin x + 2\sqrt{2}\cos x \geq \left(\sqrt{3}\right)^2$

$\Rightarrow \sin x + 2\sqrt{2}\cos x \geq 3$

$$\Rightarrow \sin\left(a + \cos^{-1}\frac{1}{3}\right) \geq 1$$

$\Rightarrow \sin\left(x + \cos^{-1}\frac{1}{3}\right) = 1$ ($\because \sin x$ cannot be greater than 1)

$$\therefore x = n\pi + (-1)^n\frac{\pi}{2} - \cos^{-1}\frac{1}{3}$$

For solution in the interval $[-2\pi, 2\pi], n = 0, 1, -1, -2$

97 **(c)**

$x^{(\log_{10} x)^2 - 3\log_{10} x + 1} > 1000 = 10^3$

$\Rightarrow [(\log_{10} x)^2 - 3\log_{10} x + 1]\log_{10} x > 3\log_{10} 10 = 3$

$\Rightarrow (\log_{10} x)^3 - 3(\log_{10} x)^2 + \log_{10} x > 3$

$\Rightarrow (\log_{10} x)(\log_{10} x - 3) + 1(\log_{10} x - 3) > 0$

$\Rightarrow (\log_{10} x - 3)(\log_{10} x + 1) > 0$

$\Rightarrow \log_{10} x - 3 > 0 \Rightarrow \log_{10} x > 3$

$\Rightarrow x > 10^3 = 1000$

$\Rightarrow x \in (1000, \infty)$

98 **(d)**

$x^{12} - x^9 + x^4 - x + 1 > 0$, three cases arise

Case I When $x \leq 0$

$x^{12} > 0, -x^9 > 0, x^4 > 0, -x > 0$

$\Rightarrow \quad x^{12} - x^9 + x^4 + x + 1 > 0, \forall\, x \leq 0$ ...(i)

Case II When $0 < x \leq 1$

$x^9 < x^4, x < 1 \Rightarrow -x^9 + x^4 > 0$ and $1 - x > 0$

$\therefore \quad x^{12} - x^9 + x^4 - x + 1 > 0, \forall\, 0 < x \leq 1$ ...(ii)

Case III When $x > 1$

$x^{12} > x^9, x^4 > x$

$\Rightarrow \quad x^{12} - x^9 + x^4 - x + 1 > 0, \forall\, x > 1$ ...(iii)

$\therefore$ From Eqs. (i), (ii) and (iii) the above equation hold for $x \in R$

100 **(a)**

We have,

$\sqrt{x+1} - \sqrt{x-1} = \sqrt{4x-1}$

$\Rightarrow x + 1 + x - 1 - 2\sqrt{x^2 - 1} = 4x - 1$

$\Rightarrow -2\sqrt{x^2 - 1} = 2x - 1$

$\Rightarrow 4(x^2 - 1) = 4x^2 - 4x + 1 \Rightarrow 4x - 5 = 0 \Rightarrow x = \frac{5}{4}$

This value of x does not satisfy the given equation.

So, the equation has no solution

101 **(d)**

The LHS of the given inequality is meaningful for $x > 0$ and $x \neq 1$

Now,

$\log_3 x - \log_x 27 < 2$

$\Rightarrow \log_3 x - 3\log_x 3 < 2$

$\Rightarrow \log_3 x - \frac{3}{\log_3 x} < 2$

$\Rightarrow \frac{(\log_3 x)^2 - 3 - 2(\log_3 x)}{\log_3 x} < 0$

$\Rightarrow \frac{(\log_3 x - 3)(\log_3 x + 1)}{(\log_3 x - 0)} < 0$

$\Rightarrow \log_3 x < -1$ or, $0 < \log_3 x < 3$

$\Rightarrow x < 3^{-1}$or, $3^0 < x < 3^3 \Rightarrow x < \frac{1}{3}$or, $1 < x < 27$

Also, $x > 0$ and $x \neq 1$

$\therefore x \in (0,1/3) \cup (1,27)$

102 **(c)**

Given inequation is $x^2 - 2x + 5 \leq 0$

$\therefore$ Roots are

$x = \frac{-2 \pm \sqrt{4-20}}{2} = \frac{2 \pm 4i}{2}$

$\therefore$ Roots are imaginary, therefore no real solutions exist

103 **(b)**

We have,

$\frac{2^{\sin^2 x} + 2^{\cos^2 x}}{2} \geq \sqrt{2^{\sin^2 x} \times 2^{\cos^2 x}}$ [Using A. M. $\geq$ G. M.]

$\Rightarrow 2^{\sin^2 x} + 2^{\cos^2 x} \geq 2\sqrt{2} \Rightarrow 2^{\sin^2 x} + 2^{\cos^2 x} \geq 2$

104 **(a)**

Given that, $a^2 + b^2 + c^2 = 1$ ...(i)

Now, $(a+b+c)^2 = a^2 + b^2 + c^2 + 2(ab + bc + ca) \geq 0$

$\Rightarrow 2(ab + bc + ca) \geq -1$ [from Eq.(i)]

$\Rightarrow ab + bc + ca \geq -\frac{1}{2}$ ...(iii)

Also, $a^2 + b^2 + c^2 - ab - bc - ca$

$= \frac{1}{2}\{(a-b)^2 + (b-c)^2 + (c-a)^2\} \geq 0$

$\Rightarrow ab + bc + ca \leq a^2 + b^2 + c^2$

$\Rightarrow ab + bc + ca \leq 1$ [from Eq.(i)] ...(iii)

From relation (ii) and (iii), we get

$-\frac{1}{2} \leq ab + bc + ca \leq 1$

105 **(b)**

We have,

$\sqrt{4x+9} - \sqrt{11x+1} = \sqrt{7x+4}$

$\Rightarrow \sqrt{4x+9} - \sqrt{7x+4} = \sqrt{11x+1}$

$\Rightarrow 4x + 9 + 7x + 4 - 2\sqrt{(4x+9)(7x+4)} = 11x + 1$

$\Rightarrow -2\sqrt{(4x+9)(7x+4)} = -12$

$\Rightarrow (4x+9)(7x+4) = 36$

$\Rightarrow 28x^2 + 79x = 0 \Rightarrow x = 0, -\frac{79}{28}$

Clearly, only $x = 0$ satisfies the given equation

106 **(a)**

Since, $-3 < x + \frac{2}{x} < 3$

$\Rightarrow -3 < \frac{(x^2+2)x}{x^2} < 3$

$\Rightarrow -3x^2 < (x^2+2)x < 3x^2 (x \neq 0)$

$\Rightarrow x(x^2 + 3x + 2) > 0$

And $x(x^2 - 3x + 2) < 0 \quad (x \neq 0)$

$\Rightarrow x(x+1)(x+2) > 0$

And $x(x-1)(x-2) < 0$

$\Rightarrow x \in (-2,-1) \cup (0,\infty)$ ...(i)

And $x \in (-\infty, 0) \cup (1,2)$...(ii)

From relations (i) and (ii), we get

$x \in (-2,-1) \cup (1,2)$

107 **(b)**

We have a, b, c are sides of a triangle

$\therefore b + c - a > 0, c + a - b > 0, a + b - c > 0$

Let $x = b + c - a, y = c + a - b, z = a + b - c$

$\Rightarrow y + z = 2a, z + x = 2b, x + y = 2c$

Now, $\frac{a}{b+c-a} + \frac{b}{c+a-b} + \frac{c}{a+b-c}$

$= \frac{y+z}{2x} + \frac{z+x}{2y} + \frac{x+y}{2z}$

$= \frac{1}{2}\left(\frac{x}{y} + \frac{y}{z} + \frac{z}{x} + \frac{x}{z} + \frac{y}{x} + \frac{z}{y}\right)$

$\geq \frac{6}{2}\left(\frac{y}{x} \cdot \frac{x}{y} \cdot \frac{y}{z} \cdot \frac{z}{y} \cdot \frac{z}{x} \cdot \frac{x}{z}\right)$ ($\because$ AM $\geq$ GM)

$= 3$

108 **(b)**

Let d_1, d_2 be the lengths of diagonals and θ be the angle between them. Then,

$$\text{Area} = \frac{1}{2}d_1 d_2 \sin\theta \Rightarrow a^2 = \frac{1}{2}d_1 d_2 \sin\theta \Rightarrow d_1 d_2 = \frac{2a^2}{\sin\theta}$$

Using A. M. $\geq$ G. M., we have

$$\frac{d_1 + d_2}{2} \geq \sqrt{d_1 d_2} \Rightarrow d_1 + d_2 \geq 2\sqrt{\frac{2a^2}{\sin\theta}} \geq 2\sqrt{2}a$$

109 **(c)**

We have, $|2x - 3| < |x + 5|$

$$\Rightarrow |2x - 3| - |x + 5| < 0$$

$$\Rightarrow \begin{cases} 3 - 2x + x + 5 < 0, x \leq -5 \\ 3 - 2x - x - 5 < 0, -5 < x \leq \frac{3}{2} \\ 2x - 3 - x - 5 < 0, x > \frac{3}{2} \end{cases}$$

$$\Rightarrow \begin{cases} x > 8, x \leq -5 \\ x > -\frac{2}{3}, -5 < x \leq \frac{3}{2} \\ x < 8, x > \frac{3}{2} \end{cases}$$

$$\Rightarrow x \in \left(-\frac{2}{3}, \frac{3}{2}\right] \cup \left(\frac{3}{2}, 8\right)$$

$$\Rightarrow x \in \left(-\frac{2}{3}, 8\right)$$

110 **(c)**

We have,

$|a_n| < 2$ i. e. $-2 < a_n < 2$

$\therefore \max(1 + a_1 x + a_2 x^2 + \cdots a_n x^n)$

$= 1 + 2|x| + 2|x|^2 + \cdots + 2|x|^n$

$= 1 + 2|x|\left\{\frac{1 - |x|^n}{1 - |x|}\right\}$

$= 1 + 2 \cdot \frac{1}{3}\left\{\frac{1 - 1/3^n}{1 - 1/3}\right\} > 0$

and,

$\min(1 + a_1 x + a_2 x^2 + \cdots + a_n x^n)$

$= 1 - 2|x| - 2|x|^2 \ldots - 2|x|^n$

$= -2[1 + |x| + |x|^2 + \cdots + |x|^n] + 3$

$= -2\left\{\frac{1 - |x|^n}{1 - |x|}\right\} + 3$

$= -2\left\{\frac{1 - 1/3^n}{1 - 1/3}\right\} + 3 > 0$

Thus, the curve $y = 1 + a_1 x + a_2 x^2 + \cdots + a_n x^n$ does not meet x-axis for $|x| < 1/3$ and $|a_n| < 2$

Hence, the equation has no real roots

112 **(c)**

Since, $|r| < 1 \Rightarrow -1 < r < 1$

Also, $a = 5(1 - r)$

$\Rightarrow 0 < a < 10$ $\left[\because \begin{matrix} \text{at } r = -1, \ a = 0 \\ \text{and at } r = 1, \ a = 0 \end{matrix}\right]$

113 **(a)**

Consider the curves $y = e^{x-8}$ and $y = 17 - 2x$. These two curves intersect at $(8, 1)$ only. Hence, the equation $e^{x-8} + 2x - 17 = 0$ has exactly one root which is equal to 8

114 **(b)**

Let $x^2 + 18x + 30 = y$. Then,

$x^2 + 18x + 30 = 2\sqrt{x^2 + 18x + 45}$

$\Rightarrow y = 2\sqrt{y + 15}$

$\Rightarrow y^2 - 4y - 60 = 0 \Rightarrow (y - 10)(y + 6) = 0 \Rightarrow y = 10$

$\therefore x^2 + 18x + 30 = y \Rightarrow x^2 + 18x + 20 = 0$

$\therefore$ Product of roots $= 20$

115 **(a)**

Since, $2 \leq |x - 3| < 4$

$\Rightarrow 2 \leq x - 3 < 4$

Or $2 \leq -(x - 3) < 4$

$\Rightarrow 5 \leq x < 7$ or $-1 \leq -x < 1$

$\Rightarrow 5 \leq x < 7$ or $-1 < x \leq 1$

$\therefore x \in (-1, 1] \cup [5, 7)$

116 **(b)**

Given that, $x = \left[\frac{a+2b}{a+b}\right]$ and $y = \frac{a}{b}$

$\therefore x = \frac{a + 2b}{a + b} = \frac{\frac{a}{b} + 2}{1 + \frac{a}{b}} = 1 + \frac{1}{\frac{a}{b} + 1}$

$\Rightarrow x = 1 + \frac{1}{y + 1}$ $\left[\because y = \frac{a}{b} \text{and} y^2 > 2 \text{ (given)}\right]$

Which shows $x^2 < 2$ $\left[\because \frac{1}{y+1} < \text{as } y > 1\right]$

117 **(c)**

Using A. M. $\geq$ G. M., we have

$5^{\sin x - 1} + 5^{-\sin x - 1} \geq 2\sqrt{5^{\sin x - 1} \times 5^{-\sin n - 1}}$

$\Rightarrow 5^{\sin x - 1} + 5^{-\sin x - 1} \geq \frac{2}{5}$

118 **(d)**

As we know, if $ax + bx + c > 0$, then $a > 0$ and $D < 0$

$\therefore (2)^2 - 4(n - 10) < 0 \Rightarrow n > 11$

119 **(a)**

Since, AM $\geq$ GM

$$\Rightarrow \frac{bcx + cay + abz}{3} \geq (a^2 b^2 c^2 \cdot xyz)^{1/3}$$

$$\Rightarrow bcx + cay + abz \geq 3\,abc \quad (\because xyz = abc)$$

120 **(b)**

Since, $\frac{(1+a_1)}{2} \geq \sqrt{1 \cdot a_1} = \sqrt{a_1}$

$\frac{(1+a_2)}{2} \geq \sqrt{1 \cdot a_2} = \sqrt{a_2}$

$\vdots \qquad \vdots$

$\frac{(1+a_n)}{2} \geq \sqrt{1 \cdot a_n} = \sqrt{a_n}$

$\Rightarrow \frac{1}{2^n}(1+a_1)(1+a_2)\ldots(1+a_n) \geq \sqrt{a_1 a_2 \ldots a_n}$
$= 1$

$\Rightarrow (1+a_1)(1+a_2)\ldots(1+a_n) \geq 2^n$

121 **(a)**

Using A. M. $\geq$ G. M., we have

$a+b \geq 2\sqrt{ab}, b+c \geq 2\sqrt{bc}$ and $c+a \geq 2\sqrt{ca}$

$\Rightarrow (a+b)(b+c)(c+a) \geq 8abc$

$\Rightarrow (2-a)(2-b)(2-c)$
$\geq 8abc \left[\because a+b+c=2 \atop \therefore b+c=2-a \text{ etc}\right]$

122 **(a)**

We have, $x^2+y^2+z^2=27$

Now,

$\frac{(x^2)^{3/2}+(y^2)^{3/2}+(z^2)^{3/2}}{3} \geq \left(\frac{x^2+y^2+z^2}{z}\right)^{3/2}$

$\Rightarrow x^3+y^3+z^3 \geq 81$

123 **(c)**

Given, $2x-7<11, 3x+4<-5$

$\Rightarrow \quad x<9, x<-3$

$\Rightarrow \quad x<-3$

$\therefore x$ lies in the interval $(-\infty, -3)$

124 **(c)**

Let

$f(x) = x^8 - x^5 - \frac{1}{x} + \frac{1}{x^4} = \frac{x^{12}-x^9-x^3+1}{x^4}$
$= \frac{(x^9-1)(x^3-1)}{x^4}$

Clearly, $f(x) \geq 0$ for all $x<0$ and it is not defined for $x=0$

For $0<x<1$, we have

$x^9-1<0$ and $x^3-1<0 \Rightarrow f(x)>0$

For $x \geq 1$, we have $x^9-1 \geq 0$ and $x^3-1 \geq 0 \Rightarrow f(x) \geq 0$

Hence, $f(x) \geq 0$ for all $x \neq 0$

125 **(c)**

We have,

$5^{(1/4)(\log_5 x)^2} \geq 5\, x^{(1/5)(\log_5 x)}$

$\frac{1}{4}(\log_5 x)^2 \log_5 5 \geq \log_5 5 + \frac{1}{5}(\log_5 x)\log_5 x$

$\Rightarrow (\log_5 x)^2 \geq 20$

$\Rightarrow (\log_5 x)^2 - \left(2\sqrt{5}\right)^2 \geq 0$

$\Rightarrow \log_5 x \leq -2\sqrt{5} \text{or}, \log_5 x \geq 2\sqrt{5}$

$\Rightarrow x \leq 5^{-2\sqrt{5}} \text{or}, x \geq 5^{2\sqrt{5}}$

$\Rightarrow x$

$\in \left(0, 5^{-2\sqrt{5}}\right]$

$\cup \left[5^{2\sqrt{5}}, \infty\right)$ [$\log_5 x$ is defined for $x>0$]

126 **(a)**

We know that,

AM $\geq$ GM

$\Rightarrow \frac{a+b}{2} \geq \sqrt{ab}$

$\Rightarrow 4 \geq \sqrt{ab}$ ($\because a+b=8$ given)

$\Rightarrow ab \leq 16$

Equality holds when number are equal. So, ab is equal to 16 when $a=4, b=4$

127 **(a)**

Curves $y=\cos x$ and $y=-|x|$ do not intersect.

So, the equation $\cos x + |x| = 0$ has no real root

128 **(d)**

Using A. M. $\geq$ G. M., we have

$\frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x} \geq 2\sqrt{\frac{\cos^3 x}{\sin x} \times \frac{\sin^3 x}{\cos x}}$ for all x
$\in (0, \pi/2)$

$\Rightarrow \frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x} \geq \sin 2x$ for all $x \in (0, \pi/2)$

$\Rightarrow \frac{\cos^3 x}{\sin x} + \frac{\sin^3 x}{\cos x} \geq 1$ for all $x \in (0, \pi/2)$

129 **(d)**

$x^2+4ax+20$

$\therefore \quad (4a)^2 - 4 \times 2 < 0$

[$\because$ if $f(x)>0$, then]

$\Rightarrow \quad 16a^2 < 8 \Rightarrow a^2 < \frac{1}{2}$

$\Rightarrow \quad -\frac{1}{\sqrt{2}} < a < \frac{1}{\sqrt{2}}$

130 **(b)**

Using A. M. $\geq$ G. M., we have

$\frac{a+b}{2} > \sqrt{ab}$ [$\because a \neq b$]

$\Rightarrow a+b > 2\sqrt{ab}$

131 **(a)**

We have,

$x^2 \cdot 2^{x+1} + 2^{|x-3|+2} = x^2 \cdot 2^{|x-3|+4} + 2^{x-1}$

Now, two cases arise

CASE I $When x \geq 3$:

In this case, we have $|x-3| = x-3$

So, the given equation reduces to

$x^2 \cdot 2^{x+1} + 2^{x-1} = x^2 \cdot 2^{x+1} + 2^{x-1}$

Which is an identity in x and hence it is true for all $x \geq 3$

CASE II $When x < 3$:

In this case, we have $|x-3| = -(x-3)$

So, the given equation reduces to

$x^2 \cdot 2^{x+1} + 2^{-x+5} = x^2 \cdot 2^{-x+7} + 2^{x-1}$

$\Rightarrow x^2 2^{x+1} - 2^{x-1} = x^2 \cdot 2^{-x+7} - 2^{-x+5}$

$\Rightarrow 2^{x-1}(4x^2-1) = 2^{-x+5}(4x^2-1)$

$\Rightarrow 2^{2x}(4x^2-1) = 2^6(4x^2-1)$

$\Rightarrow (2^{2x} - 2^6)(4x^2-1) = 0$

$\Rightarrow 2x = 6$ or, $4x^2 - 1 = 0$

$\Rightarrow x = 3$ or, $x = \pm\frac{1}{2}$

But, $x < 3$. Therfore, $x = \pm\frac{1}{2}$

Hence, the given equation has no negative integral root

132 **(b)**

We have,

$2^{\sin^2 x} \cdot 3^{\cos^2 y} \cdot 4^{\sin^2 z} \cdot 5^{\cos^2 \omega} \geq 120$

$\Rightarrow 2^{\sin^2 x} \cdot 3^{\cos^2 y} \cdot 4^{\sin^2 z} \cdot 5^{\cos^2 \omega} \geq 2 \times 3 \times 4 \times 5$

$\Rightarrow \sin^2 x \log 2 + \cos^2 y \log_3 + \sin^2 z \log 4 + \cos^2 \omega \log 5 \geq \log 2 + \log 3 + \log 4 + \log 5$

$\Rightarrow \cos^2 x \log 2 + \sin^2 y \log 3 + \cos^2 z \log 4 + \sin^2 \omega \log 5 \leq 0$

$\Rightarrow \cos^2 x = 0, \sin^2 y = 0, \cos^2 z = 0$ and $\sin^2 \omega = 0$

$\Rightarrow x = m\pi \pm \frac{\pi}{2}, m \in Z; y = n\pi, n \in Z$

$z = r\pi \pm \frac{\pi}{2}, r \in Z;\ \omega = t\,\pi, t \in Z$

But, $x, y, z, \omega \in [0,10]$

$\therefore x = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, y = 0, \pi, 2\pi, 3\pi, z = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}$

and $\omega = 0, \pi, 2\pi, 3\pi$

Hence, the number of ordered 4-tuples is $3 \times 4 \times 3 \times 4 = 144$

133 **(d)**

We have,

$\log_b a + \log_a b + \log_d c + \log_c d$

$= \left(\log_b a + \frac{1}{\log_b a}\right) + \left(\log_d c + \frac{1}{\log_d c}\right) \geq 2 + 2 = 4$

134 **(c)**

We have,

$\log_{1/3}(2^{x+2} - 4^x) \geq -2$

$\Rightarrow 2^{x+2} - 4^x \leq \left(\frac{1}{3}\right)^{-2}$ and $2^{x+2} - 4^x > 0$

$\Rightarrow 4(2^x) - (2^x)^2 \leq 9$ and $2^x(2^2 - 2^x) > 0$

$\Rightarrow (2^x)^2 - 4(2^x) + 9 \geq 0$ and $2^x < 2^2$

$\Rightarrow x < 2$ $[\because (2^x)^2 - 4(2^x) + 9 > 0$ for all $x \in R]$

$\Rightarrow x \in (-\infty, 2)$

135 **(c)**

Given, $\frac{2x}{(2x+1)(x+2)} - \frac{1}{(x+1)} > 0$

$\Rightarrow \frac{-3x-2}{(x+1)(x+2)(2x+1)} > 0$

Equating each factor equal to 0, we have

$x = -2, -1, -\frac{2}{3}, -\frac{1}{2}$

It is clear $-\frac{2}{3} < x < -\frac{1}{2}$ or $-2 < x < -1$

136 **(b)**

We observe that the curves $y = \log_2 x$ and $y = -|x|$ intersect at exactly one point. So, the equation $\log_2 x + |x| = 0$ has exactly one real root



137 **(a)**

Using A. M. $\geq$ G. M., we have

$\frac{bcx + cay + abz}{3} \geq (bcx \times cay \times abz)^{1/3}$

$\Rightarrow bcx + cay + abz \geq 3(a^2b^2c^2 \times xyz)^{1/3}$

$\Rightarrow bcx + cay + abz \geq 3abc$ $[\because xyz = abc]$

138 **(a)**

We have,

$\sqrt{3x^2 + 6x + 7} + \sqrt{5x^2 + 10x + 14} \leq 4 - 2x - x^2$

$\Rightarrow \sqrt{3(x+1)^2 + 4} + \sqrt{5(x+1)^2 + 9} \leq (x+1)^2 + 5$

Clearly, LHS ≥ 5 and LHS ≤ 5

So, the inequation holds when each side is equal to 5

This is true when $x = -1$

Hence, the given inequation has exactly one solution

139 **(b)**

Let $a_1, a_2, a_3, \ldots, a_n$ be the lengths of n parts of the stick. Then,

$a_1 + a_2 + a_3 + \cdots + a_n = 20$ and $a_1 a_2 a_3 \ldots a_n > 1$

Now, A. M. ≥ G. M.

$\Rightarrow \frac{a_1 + a_2 + \cdots + a_n}{a_n} \geq (a_1 a_2 \ldots a_n)^{1/n}$

$\Rightarrow \frac{20}{n} > 1 \Rightarrow n < 20$

$\therefore$ Maximum possible value of n is 19

140 **(b)**

$\because$ AM $> GM$

$$\frac{(a+b-c)+(b+c-a)}{2} > \sqrt{(a+b-c)(b+c-a)}$$

$\Rightarrow b > \sqrt{(a+b-c)(b+c-a)}$...(i)

Similarly, $\frac{(b+c-a)+(c+a-b)}{2} > \sqrt{(b+c-a)(c+a-b)}$

$\Rightarrow c > \sqrt{(b+c-a)(c+a-b)}$...(ii)

and $\frac{(c+a-b)+(a+b-c)}{2} > \sqrt{(c+a-b)(a+b-c)}$

$\Rightarrow a > \sqrt{(c+a-b)(a+b-c)}$...(iii)

On multiplying relations (i), (ii) and (iii), we get

$abc > (a+b-c)(b+c-a)(c+a-b)$

$\Rightarrow (a+b-c)(b+c-a)(c+a-b) - abc < 0$

141 **(c)**

We have,

$\log_{16} x^3 + \left(\log_2 \sqrt{x}\right)^2 < 1$

$\Rightarrow \frac{3}{4}\log_2 x + \frac{1}{4}(\log_2 x)^2 < 1$

$\Rightarrow (\log_2 x)^2 + 3\log_2 x - 4 < 0$

$\Rightarrow (\log_2 x + 4)(\log_2 x - 1) < 0$

$\Rightarrow -4 < \log_2 x < 1 \Rightarrow 2^{-4} < x < 2 \Rightarrow x \in (1/16, 2)$

Also, LHS of the given inequality is defined fro $x > 0$

Hence, $x \in (1/16, 2)$

142 **(b)**

Since, $\sin x \leq \cos^2 x$, becuase $\cos x$ must be a positive proper fraction

$\sin^2 x + \sin x - 1 \leq 0$

Or $\left(\sin x + \frac{1}{2}\right)^2 - \frac{5}{4} \leq 0$

From the definition of logarithm

$\sin x > 0, \cos x > 0, \cos x \neq 1$

$\therefore \sin x + \frac{1}{2} \leq \frac{\sqrt{5}}{2}$,

$\Rightarrow 0 < \sin x \leq \frac{\sqrt{5}-1}{2}$

143 **(d)**

If $f(x) = x^2 + 2bx + 2c^2$ and $g(x) = -x^2 - 2cx + b^2$

Then, $f(x)$ is minimum and $g(x)$ is maximum at $f(x) = \frac{-D}{4a}$, $\left(\because x = -\frac{-b}{a} \text{and } f(x) = \frac{-D}{4a}\right)$

$\therefore \min\{f(x)\} = \frac{-(4b^2 - 8c^2)}{4} = (2c^2 - b^2)$

And $\max\{g(x)\} = -\frac{(4c^2+4b^2)}{4(-1)} = (b^2 + c^2)$

Since, $\min f(x) > \max g(x) \Rightarrow 2c^2 - b^2 > b^2 + c^2$

$\Rightarrow \quad c^2 > 2b^2 \quad \Rightarrow \quad |c| > \sqrt{2}|b|$

144 **(a)**

We have, $[x] + (x) = 5$

If $x \leq 2$, then $[x] + (x) \leq 2 + 2 < 5$

If $x \geq 3$, then $[x] + (x) \geq 3 + 3 > 5$

If $2 < x < 3$, then $[x] + (x) = 2 + 3 = 5$

Hence, the solution set is $(2, 3)$

145 **(a)**

We have,

$|x - x^2 - 1| = |2x - 3 - x^2|$

$\Rightarrow |x^2 - x + 1| = |x^2 - 2x + 3|$

$\Rightarrow x^2 - x + 1$

$= x^2 - 2x + 3 \quad \left[\begin{matrix}\because x^2 - x + 1 > 0 \\ \text{and } x^2 - 2x + 3 > 0 \text{ for all } x\end{matrix}\right]$

$\Rightarrow x = 2$

146 **(a)**

Given, $\frac{|x-1|}{x+2} - 1 < 0$

Case I When $x < 1, |x - 1| = 1 - x$

$\therefore \frac{1-x}{x+2} - 1 < 0 \Rightarrow \frac{-2x-1}{x+2} < 0$

$\Rightarrow \frac{2x+1}{x+2} > 0 \Rightarrow x < -2 \text{ or } x > -\frac{1}{2}$

But $x < 1$

$\therefore x \in (-\infty, -2) \cup \left(-\frac{1}{2}, 1\right)$

Case II When $x \geq 1, |x - 1| = x - 1$

$\therefore \frac{x-1}{x+2} - 1 < 0 \Rightarrow -\frac{3}{x+2} < 0$

$\Rightarrow \frac{3}{x+2} > 0$

$\Rightarrow \quad x > -2$

But $x \geq 1$

$\therefore \quad x \geq 1, ie, x \in [1, \infty)$ ...(iii)

$\therefore$ From Eqs. (i) and (ii), we get

$x \in (-\infty, -2) \cup \left(-\frac{1}{2}, \infty\right)$

147 **(d)**

Given that, $\frac{x+2}{x^2+1} > \frac{1}{2}$

$\Rightarrow \quad x^2 - 2x - 3 < 0$

$\Rightarrow \quad (x-3)(x+1) < 0$

$\Rightarrow \ -1 < x < 3$

The integer value of x are 0, 1, 2

$\therefore$ The number of integral solutions are 3

148 **(b)**

$\because \tan\left(x + \frac{\pi}{3}\right) \geq 1 \Rightarrow \frac{\pi}{4} \leq x + \frac{\pi}{3} < \frac{\pi}{2}$

$$\Rightarrow -\frac{\pi}{12} \leq x < \frac{\pi}{6}$$

$$\Rightarrow n\pi - \frac{\pi}{12} \leq x \leq n\pi + \frac{\pi}{6}$$

149 **(c)**

We have,

$$\frac{1 + (\log_a x)^2}{1 + \log_a x} > 1$$

$$\Rightarrow \frac{1 + (\log_a x)^2}{1 + \log_a x} - 1 > 0$$

$$\Rightarrow \frac{(\log_a x)(\log_a x - 1)}{(1 + \log_a x)} > 0$$

$\Rightarrow -1 < \log_a x < 0$ or, $\log_a x > 1$

$\Rightarrow a^{-1} > x > a^0$ or, $x < a [\because 0 < a < 1]$

$\Rightarrow x \in (1, 1/a) \cup (0, a) [\because a > 0]$

150 **(b)**

We have,

$$x = \frac{y+2}{y+1}$$

$$\Rightarrow y = \frac{2-x}{x-1}$$

$$\Rightarrow \left(\frac{2-x}{x-1}\right)^2 > 2 \qquad [\because y^2 > 2]$$

$\Rightarrow (2-x)^2 > 2(x-1)^2 \Rightarrow x^2 < 2$

151 **(a)**

Let $x = y - \frac{3\pi}{4}$. Then,

$$\sin x = -\left(\frac{\cos y + \sin y}{\sqrt{2}}\right) \text{ and } \cos x = -\left(\frac{\cos y - \sin y}{\sqrt{2}}\right)$$

$\Rightarrow \sin x + \cos x = -\sqrt{2}\cos y$ and $\sin x \cos x = \frac{1}{2}(2\cos^2 y - 1)$

Now,

$$|\sin x + \cos x + \tan x + \sec x + \operatorname{cosec} x + \cot x|$$

$$= |(\sin x + \cos x) + (\tan x + \cot x) + (\sec x + \operatorname{cosec} x)|$$

$$= \left|(\sin x + \cos x) + \frac{1}{\sin x \cos x} + \frac{\sin x + \cos x}{\sin x \cos x}\right|$$

$$= \left|(\sin x + \cos x)\left(1 + \frac{1}{\sin x \cos x}\right) + \frac{1}{\sin x \cos x}\right|$$

$$= \left|-\sqrt{2}\cos y\left(1 + \frac{2}{2\cos^2 y - 1}\right) + \frac{2}{2\cos^2 y - 1}\right|$$

$$= \left|-\sqrt{2}\cos y - \frac{2\left(\sqrt{2}\cos y - 1\right)}{2\cos^2 y - 1}\right|$$

$$= \left|-\sqrt{2}\cos y - \frac{2}{\sqrt{2}\cos y + 1}\right|$$

$$= \left|\sqrt{2}\cos y + \frac{2}{\sqrt{2}\cos y + 1}\right|$$

$$= \left|\lambda + \frac{2}{\lambda + 1}\right|, \text{where } \lambda = \sqrt{2}\cos y$$

$$= \left|(\lambda + 1) + \frac{2}{\lambda + 1} - 1\right| \geq \left|(\lambda + 1) + \frac{2}{(\lambda + 1)}\right| - 1$$

$$\geq 2\sqrt{(\lambda + 1) \times \frac{2}{(\lambda + 1)}} - 1 = 2\sqrt{2} - 1 \text{ [Using AM} \geq \text{GM]}$$

152 **(b)**

$$\frac{9 \cdot 3^{2x} - 6 \cdot 3^x + 4}{9 \cdot 3^{2x} + 6 \cdot 3^x + 4} = \frac{\left(3^{(x+1)}\right)^2 - 2(3^{(x+1)}) + 4}{(3^{(x+1)})^2 + 2(3^{(x+1)}) + 4}$$

$$= \frac{t^2 - 2t + 4}{t^2 + 2t + 4} \quad (\text{where } t = 3^{x+1}) \quad \ldots(\text{i})$$

Since, $\frac{1}{3} < \frac{x^2 - 2x + 4}{x^2 + 2x + 4} < 3$

$\therefore$ From Eq.(i), the given expression lies between 1/3 and 3

153 **(c)**

Using A. M. $\geq$ G. M., we have

$4^x + 4^{1-x} \geq 2\sqrt{4^x \times 4^{1-x}} \Rightarrow 4^x + 4^{1-x} \geq 4$

154 **(b)**

We have,

$3^{-|x|} - 2^{|x|} = 0 \Rightarrow 3^{-|x|} = 2^{|x|} \Rightarrow 6^{|x|} = 1 \Rightarrow x = 0$

156 **(d)**

We have,

$P \leq (x_1 + x_2 + \cdots + x_{n-1})(x_2 + x_4 + x_6 + \cdots + x_n)$

$$\Rightarrow P \leq \frac{1}{4}(x_1 + x_2 + \cdots + x_n)^2 = \frac{1}{4}$$

157 **(b)**

We observe that $y = e^{-x}$ and $y = x$ intersect at exactly one point. So, the equation $e^{-x} = x$ has exactly one real root:



158 **(b)**

We have,

$3^x + 3^{1-x} - 4 < 0$

$\Rightarrow (3^x)^2 - 4(3^x) + 3 < 0$

$\Rightarrow (3^x - 1)(3^x - 3) < 0$

$\Rightarrow 1 < 3^x < 3 \Rightarrow 0 < x < 1 \Rightarrow x \in (0, 1)$

160 **(c)**

We know that $x + \frac{1}{x} \geq 2$ for all $x > 0$

$\therefore \sin^5\theta + \text{cosec}^5\theta \geq 2$ for $0 < \theta < \pi$

Hence, the minimum value of $\sin^5\theta + \text{cosec}^5\theta$ is 2

1)	a	2)	c	3)	d	4)	b	85)	c	86)	c	87)	d	88)	c
5)	c	6)	a	7)	c	8)	c	89)	c	90)	b	91)	b	92)	d
9)	a	10)	c	11)	b	12)	c	93)	c	94)	a	95)	d	96)	d
13)	b	14)	b	15)	d	16)	c	97)	c	98)	d	99)	d	100)	a
17)	b	18)	c	19)	b	20)	d	101)	d	102)	c	103)	b	104)	a
21)	b	22)	a	23)	a	24)	b	105)	b	106)	a	107)	b	108)	b
25)	b	26)	c	27)	a	28)	b	109)	c	110)	c	111)	d	112)	c
29)	a	30)	b	31)	a	32)	b	113)	a	114)	b	115)	a	116)	b
33)	a	34)	d	35)	c	36)	b	117)	c	118)	d	119)	a	120)	b
37)	a	38)	a	39)	b	40)	b	121)	a	122)	a	123)	c	124)	c
41)	c	42)	c	43)	a	44)	a	125)	c	126)	a	127)	a	128)	d
45)	a	46)	c	47)	b	48)	a	129)	d	130)	b	131)	a	132)	b
49)	a	50)	c	51)	a	52)	d	133)	d	134)	c	135)	c	136)	b
53)	d	54)	d	55)	c	56)	d	137)	a	138)	a	139)	b	140)	b
57)	a	58)	d	59)	c	60)	a	141)	c	142)	b	143)	d	144)	a
61)	b	62)	c	63)	c	64)	c	145)	a	146)	a	147)	d	148)	b
65)	b	66)	d	67)	a	68)	a	149)	c	150)	b	151)	a	152)	b
69)	c	70)	d	71)	a	72)	a	153)	c	154)	b	155)	b	156)	d
73)	c	74)	a	75)	c	76)	c	157)	b	158)	b	159)	b	160)	c
77)	d	78)	d	79)	a	80)	d								
81)	b	82)	a	83)	b	84)	a								

Click here to Download more Question Bank for Additional Chapters.

Dear Teachers and Students,

Join School of Educators' exclusive WhatsApp, Telegram, and Signal groups for FREE access to a vast range of educational resources designed to help you achieve 100/100 in exams! Separate groups for teachers and students are available, packed with valuable content to boost your performance.

Additionally, benefit from expert tips, practical advice, and study hacks designed to enhance performance in both CBSE exams and competitive entrance tests.

Don’t miss out—join today and take the first step toward academic excellence!


Join the Teachers and Students Group by Clicking the Link Below
What’s app (Teachers)
Telegram
Signal
What’s app (Students)
Join Youtube
CBSE Skill courses
Learn more about groups www.schoolofeducators.com

SCHOOL OF EDUCATORS

SOE

Learn | Unlearn | Relearn

JOIN OUR WHATSAPP GROUPS

FOR FREE EDUCATIONAL RESOURCES

JOIN SCHOOL OF EDUCATORS WHATSAPP GROUPS FOR FREE EDUCATIONAL RESOURCES

We are thrilled to introduce the School of Educators WhatsApp Group, a platform designed exclusively for educators to enhance your teaching & Learning experience and learning outcomes. Here are some of the key benefits you can expect from joining our group:

BENEFITS OF SOE WHATSAPP GROUPS

- **Abundance of Content:** Members gain access to an extensive repository of educational materials tailored to their class level. This includes various formats such as PDFs, Word files, PowerPoint presentations, lesson plans, worksheets, practical tips, viva questions, reference books, smart content, curriculum details, syllabus, marking schemes, exam patterns, and blueprints. This rich assortment of resources enhances teaching and learning experiences.

- **Immediate Doubt Resolution:** The group facilitates quick clarification of doubts. Members can seek assistance by sending messages, and experts promptly respond to queries. This real-time interaction fosters a supportive learning environment where educators and students can exchange knowledge and address concerns effectively.

- **Access to Previous Years' Question Papers and Topper Answers:** The group provides access to previous years' question papers (PYQ) and exemplary answer scripts of toppers. This resource is invaluable for exam preparation, allowing individuals to familiarize themselves with the exam format, gain insights into scoring techniques, and enhance their performance in assessments.

- **Free and Unlimited Resources:** Members enjoy the benefit of accessing an array of educational resources without any cost restrictions. Whether its study materials, teaching aids, or assessment tools, the group offers an abundance of resources tailored to individual needs. This accessibility ensures that educators and students have ample support in their academic endeavors without financial constraints.

- **Instant Access to Educational Content:** SOE WhatsApp groups are a platform where teachers can access a wide range of educational content instantly. This includes study materials, notes, sample papers, reference materials, and relevant links shared by group members and moderators.

- **Timely Updates and Reminders:** SOE WhatsApp groups serve as a source of timely updates and reminders about important dates, exam schedules, syllabus changes, and academic events. Teachers can stay informed and well-prepared for upcoming assessments and activities.

- **Interactive Learning Environment:** Teachers can engage in discussions, ask questions, and seek clarifications within the group, creating an interactive learning environment. This fosters collaboration, peer learning, and knowledge sharing among group members, enhancing understanding and retention of concepts.

- **Access to Expert Guidance:** SOE WhatsApp groups are moderated by subject matter experts, teachers, or experienced educators can benefit from their guidance, expertise, and insights on various academic topics, exam strategies, and study techniques.

Join the School of Educators WhatsApp Group today and unlock a world of resources, support, and collaboration to take your teaching to new heights. To join, simply click on the group links provided below or send a message to +91-95208-77777 expressing your interest.

Together, let's empower ourselves & Our Students and inspire the next generation of learners.

Best Regards,
Team
School of Educators

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Teachers Only)

Class 1

Class 2

Class 3

Class 4

Class 5

Class 6

Class 7

Class 8

Class 9

Class 10

Class 11 (Science)

Class 11 (Humanities)

Class 11 (Commerce)

Class 12 (Science)

Class 12 (Humanities)

Class 12 (Commerce)

Kindergarten

Subject Wise Secondary and Senior Secondary Groups (IX & X For Teachers Only)

Secondary Groups (IX & X)

Click Here to Join
SST

Click Here to Join
Mathematics

Click Here to Join
Science

Click Here to Join
English

Click Here to Join
Hindi-A

Click Here to Join
IT Code-402

Click Here to Join
Hindi-B

Click Here to Join
Artificial Intelligence

Senior Secondary Groups (XI & XII For Teachers Only)

Click Here to Join
Physics

Click Here to Join
Chemistry

Click Here to Join
English

Click Here to Join
Mathematics

Click Here to Join
Biology

Click Here to Join
Accountancy

Click Here to Join
Economics

Click Here to Join
BST

Click Here to Join
History

Click Here to Join

Geography

Click Here to Join

Sociology

Click Here to Join

Hindi Elective

Click Here to Join

Hindi Core

Click Here to Join

Home Science

Click Here to Join

Sanskrit

Click Here to Join

Psychology

Click Here to Join

Political Science

Click Here to Join

Painting

Click Here to Join

Vocal Music

Click Here to Join

Comp. Science

Click Here to Join

IP

Click Here to Join

Physical Education

Click Here to Join

APP. Mathematics

Click Here to Join

Legal Studies

Click Here to Join

Entrepreneurship

Click Here to Join

French

Click Here to Join

IT

Click Here to Join

Artificial Intelligence

Other Important Groups (For Teachers & Principal's)

Click Here to Join

Principal's Group

Click Here to Join

Teachers Jobs

Click Here to Join

IIT/NEET

Join School of Educators WhatsApp Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class WhatsApp Group.

Kindergarten to Class XII (For Students Only)

Class 1

Class 2

Class 3

Class 4

Class 5

Class 6

Class 7

Class 8

Class 9

Class 10

Class 11 (Science)

Class 11 (Humanities)

Class 11 (Commerce)

Class 12 (Science)

Class 12 (Humanities)

Class 12 (Commerce)

Artificial Intelligence
(VI TO VIII)

Subject Wise Secondary and Senior Secondary Groups (IX & X For Students Only)

Secondary Groups (IX & X)

SST

Mathematics

Science

English

Hindi

IT Code

Artificial Intelligence

Senior Secondary Groups (XI & XII For Students Only)

Physics

Chemistry

English

Mathematics

Biology

Accountancy

Economics

BST

History

Click Here to Join

Geography

Click Here to Join

Sociology

Click Here to Join

Hindi Elective

Click Here to Join

Hindi Core

Click Here to Join

Home Science

Click Here to Join

Sanskrit

Click Here to Join

Psychology

Click Here to Join

Political Science

Click Here to Join

Painting

Click Here to Join

Music

Click Here to Join

Comp. Science

Click Here to Join

IP

Click Here to Join

Physical Education

Click Here to Join

APP. Mathematics

Click Here to Join

Legal Studies

Click Here to Join

Entrepreneurship

Click Here to Join

French

Click Here to Join

IT

Click Here to Join

AI

Click Here to Join

IIT/NEET

Click Here to Join

CUET

Groups Rules & Regulations:

To maximize the benefits of these WhatsApp groups, follow these guidelines:

1. Share your valuable resources with the group.
2. Help your fellow educators by answering their queries.
3. Watch and engage with shared videos in the group.
4. Distribute WhatsApp group resources among your students.
5. Encourage your colleagues to join these groups.

Additional notes:

1. Avoid posting messages between 9 PM and 7 AM.
2. After sharing resources with students, consider deleting outdated data if necessary.
3. It's a NO Nuisance groups, single nuisance and you will be removed.
   - No introductions.
   - No greetings or wish messages.
   - No personal chats or messages.
   - No spam. Or voice calls
   - Share and seek learning resources only.

Please only share and request learning resources. For assistance, contact the helpline via WhatsApp: +91-95208-77777.

Join Premium WhatsApp Groups
Ultimate Educational Resources!!

Join our premium groups and just Rs. 1000 and gain access to all our exclusive materials for the entire academic year. Whether you're a student in Class IX, X, XI, or XII, or a teacher for these grades, Artham Resources provides the ultimate tools to enhance learning. Pay now to delve into a world of premium educational content!

Click here for more details

Click Here to Join

Class 9

Click Here to Join

Class 10

Click Here to Join

Class 11

Click Here to Join

Class 12

📣 Don't Miss Out! Elevate your academic journey with top-notch study materials and secure your path to top scores! Revolutionize your study routine and reach your academic goals with our comprehensive resources. Join now and set yourself up for success! 📚🌟

Best Wishes,

Team
School of Educators & Artham Resources

SKILL MODULES BEING OFFERED IN MIDDLE SCHOOL

Artificial Intelligence

Beauty & Wellness

Design Thinking & Innovation

Financial Literacy

Handicrafts

Information Technology

Marketing/Commercial Application



Mass Media - Being Media Literate

Travel & Tourism

Coding



Data Science (Class VIII only)



Augmented Reality / Virtual Reality

Digital Citizenship

Life Cycle of Medicine & Vaccine

Things you should know about keeping Medicines at home



What to do when Doctor is not around



Humanity & Covid-19

Blue Pottery



Pottery

Block Printing



Food



Food Preservation



Baking

Herbal Heritage

Khadi



Mask Making

Mass Media

Making of a Graphic Novel



Kashmiri Embroidery

Embroidery

Rockets

Satellites

Application of Satellites



Photography

SKILL SUBJECTS AT SECONDARY LEVEL (CLASSES IX – X)

Retail

Information Technology



Security

Automotive

Introduction To Financial Markets

Introduction To Tourism

Beauty & Wellness

Agriculture

Food Production

Front Office Operations

Banking & Insurance

Marketing & Sales

Health Care

Apparel

Multi Media

Multi Skill Foundation Course

Artificial Intelligence

Physical Activity Trainer

Data Science

Electronics & Hardware (NEW)

Foundation Skills For Sciences (Pharmaceutical & Biotechnology)(NEW)

Design Thinking & Innovation (NEW)

SKILL SUBJECTS AT SR. SEC. LEVEL (CLASSES XI – XII)

Retail

InformationTechnology

Web Application

Automotive

Financial Markets Management

Tourism

Beauty & Wellness

Agriculture

Food Production

Front Office Operations

Banking

Marketing

Health Care

Insurance

Horticulture



Typography & Comp. Application

Geospatial Technology



Electrical Technology

Electronic Technology



Multi-Media

Taxation

Cost Accounting



Office Procedures & Practices



Shorthand (English)



Shorthand (Hindi)



Air-Conditioning & Refrigeration



Medical Diagnostics

Textile Design

Design



Salesmanship



Business Administration



Food Nutrition & Dietetics



Mass Media Studies

Library & Information Science

Fashion Studies



Applied Mathematics

Yoga



Early Childhood Care & Education

Artificial Intelligence

Data Science

Physical Activity Trainer(new)



Land Transportation Associate (NEW)

Installation Technician Computing and Peripherals
Sector: Electronics and Hardware
CENTRAL BOARD OF SECONDARY EDUCATION

Electronics & Hardware (NEW)

Design Thinking & Innovation (NEW)

Join School of Educators Signal Groups

You will get Pre- Board Papers PDF, Word file, PPT, Lesson Plan, Worksheet, practical tips and Viva questions, reference books, smart content, curriculum, syllabus, marking scheme, toppers answer scripts, revised exam pattern, revised syllabus, Blue Print etc. here . Join Your Subject / Class signal Group.

Kindergarten to Class XII

Click Here to Join
Class 1

Click Here to Join
Class 2

Click Here to Join
Class 3

Click Here to Join
Class 4

Click Here to Join
Class 5

Click Here to Join
Class 6

Click Here to Join
Class 7

Click Here to Join
Class 8

Click Here to Join
Class 9

Click Here to Join
Class 10

Click Here to Join
Class 11 (Science)

Click Here to Join
Class 11 (Humanities)

Click Here to Join
Class 11 (Commerce)

Click Here to Join
Class 12 (Science)

Click Here to Join
Class 12 (Humanities)

Click Here to Join
Class 12 (Commerce)

Click Here to Join
Kindergarten

Click Here to Join
Artifical intelligence

Subject Wise Secondary and Senior Secondary Groups IX & X

Secondary Groups (IX & X)

SST

Mathematics

Science

English

Hindi-A

Hindi-B

IT Code-402

IT

Artifical intelligence

Senior Secondary Groups XI & XII

Physics

Chemistry

English

Mathematics

Biology

Accountancy

Economics

BST

History

Click Here to Join

Geography

Click Here to Join

Sociology

Click Here to Join

Hindi Elective

Click Here to Join

Hindi Core

Click Here to Join

Home Science

Click Here to Join

Sanskrit

Click Here to Join

Psychology

Click Here to Join

Political Science

Click Here to Join

Painting

Click Here to Join

Vocal Music

Click Here to Join

Comp. Science

Click Here to Join

IP

Click Here to Join

Physical Education

Click Here to Join

APP. Mathematics

Click Here to Join

Legal Studies

Click Here to Join

Entrepreneurship

Click Here to Join

French

Click Here to Join

IIT/NEET

Click Here to Join

Artifical intelligence

Click Here to Join

CUET

Join School of Educators CBSE Telegram Groups

Join us on Telegram

Kindergarten

Join us on Telegram

All classes

Join us on Telegram

Class 1

Join us on Telegram

Class 2

Join us on Telegram

Class 3

Join us on Telegram

Class 4

Join us on Telegram

Class 5

Join us on Telegram

Class 6

Join us on Telegram

Class 7

Join us on Telegram

Class 8

Join us on Telegram

Class 9

Join us on Telegram

Class 10

Join us on Telegram

Class 11 (Sci)

Join us on Telegram

Class 11 (Com)

Join us on Telegram

Class 11 (Hum)

Join us on Telegram

Class 12 (Sci)

Join us on Telegram

Class 12 (Com)

Join us on Telegram

Class 12 (Hum)

Join us on Telegram

JEE/NEET

Join us on Telegram

CUET

Join us on Telegram

NDA, OLYMPIAD, NTSE

Join us on Telegram

Principal Professional Group

Join us on Telegram

Teachers Professional Group

Join us on Telegram

Project File Group

Join School of Educators ICSE Telegram Groups



Kindergarten



Class 1



Class 2



Class 3



Class 4



Class 5

Class 6



Class 7



Class 8

Class 9

Class 10

Class 11 (Sci)

Class 11 (Com)

Class 11 (Hum)

Class 12 (Sci)

Class 12 (Com)

Class 12 (Hum)

ARTHAM
RESOURCE MATERIAL

SCHOOL OF EDUCATORS
SOE
Learn / Unlearn / Relearn